

## A Clinical Study of Ectopic Pregnancy

Sreelatha B\*, Satyaprabha S  
and Prasanna K

### Abstract

**Background:** Ectopic pregnancy is not an uncommon condition and is potentially a life-threatening emergency. There is a rising trend in incidence which necessitates the gynecologists to have comprehensive knowledge about the disease.

**Aim of the study:** To determine the incidence, risk factors, clinical features and management.

**Materials and Methods:** This study is conducted in Department of Obstetrics and Gynecology of PIMS, Karimnagar. It is a retrospective descriptive study on patients who presented to our department with clinical features of ectopic pregnancy and later confirmed with intraoperative findings or ultrasound, in the 3-year period from January 2017 to January 2020.

**Results:** A total of 34 patients were diagnosed to have ectopic pregnancy with an incidence of 1 in 114 deliveries. The cause of ectopic pregnancy was dilatation and curettage in 11.8% and PID, infertility, prior tubal surgery, previous abortions and OCP usage each contributes by 5.9%. Surgical treatment with salpingectomy was done in majority of the cases and medical management in 3 patients. Blood transfusions were given in 38.2% of cases. Postoperative period is uneventful in all the cases.

**Conclusion:** Ectopic pregnancy is leading cause of maternal mortality in first trimester. In spite of advanced diagnostic techniques, it poses great diagnostic difficulties due to varied signs and symptoms. Previous tubal surgery pelvic inflammatory disease and infertility are the risk factors of tubal pregnancy.

**Keywords:** Ectopic pregnancy; Salpingectomy; Ovarian pregnancy

Department of Obstetrics and Gynecology,  
Prathima Institute of Medical Sciences  
(PIMS), Nagunoor, Karimnagar, Telangana,  
India

\*Corresponding author: Sreelatha B

✉ sreelathareddy1991@gmail.com

Department of Obstetrics and Gynecology,  
Prathima Institute of Medical Sciences,  
Nagunoor, Karimnagar, Telangana, India.

Tel: +91-8374883436

**Citation:** Sreelatha B, Satyaprabha S,  
Prasanna K (2021) A Clinical Study of Ectopic  
Pregnancy. Gynecol Obstet Case Rep Vol.7  
No.6:142

**Received:** May 25, 2021; **Accepted:** June 17, 2021; **Published:** June 24, 2021

### Introduction

Gynecologists should have comprehensive knowledge about the clinical presentation and risk factors of EP. If suspected clinically, encouraging women to undergo early ultrasonography allows early detection of EP and can be managed medically or fertility sparing surgical procedures. In Ectopic Pregnancy (EP) gestational sac is implanted outside the endometrial lining. The incidence of EP is around 1-2% in most hospital based studies [1-5] and has been on the rise during the last few decades.

Classic triad of amenorrhea, abdominal pain and vaginal bleeding is may not be seen in majority of cases. Women may present with non-specific symptoms, unaware of an ongoing pregnancy or even present with hemodynamic shock. The contribution of EP to the maternal mortality rates in developing countries including India is not precisely known, with data from few

studies indicating 3.5%-7.1% maternal deaths due to EP [6,7]. An accurate history and physical examination and its correlation with diagnostic techniques is important for diagnosis and management. Immediate intervention is required to prevent maternal morbidity and mortality.

### Materials and Methods

This study was conducted at Prathima Institute of Medical Sciences, a tertiary care teaching hospital in Karimnagar, South India. The study was approved by the Institute Ethics Committee. It is a retrospective descriptive study on patients who presented to our department with clinical features of ectopic pregnancy and later confirmed with intraoperative findings or ultrasound, in the 3-year period from January 2017 to January 2020. The case records of patients diagnosed with EP retrieved from the medical records department.

Patient characteristics like age, parity, presenting symptoms, clinical signs, examination findings were noted. Mode of diagnosis, management modality, complications and need for blood transfusion were also recorded. The primary outcome measures studied were incidence of EP, their risk factors, mortality and morbidity in these women.

## Results

There were 34 ectopic pregnancies in the study period. Incidence of EP is 1 in 114 deliveries (0.87%). It was more common in the age group of 21 to 30 years (70.6%) with minimum age at diagnosis made was 19 years and maximum age at diagnosis made was 38 years. 9 out of 34 patients were primigravida and rest were multiparous women. Previous C section history seen in more than half (56%) of patients. History of abortion in 6 patients (17.7%). There was previous history of ectopic pregnancy in two patients. Two patients had Infertility and took treatment for same. No risk factor was identified in one patient (Table 1).

### Presenting symptoms

Abdominal pain was the predominant symptom followed by Amenorrhea and bleeding. Nearly 1/4<sup>th</sup> of patients presented with nonspecific symptoms. On General physical examination Pallor was seen in 76.5% and fever in 5.9% patients. Two patients were in mild shock.

Abdominal palpation shows Tenderness in 27 patients and Guarding in 5 patients. Abdominal distension was seen in 5 cases. More than half (58.5%) of patients had vaginal bleed on per speculum exam. Cervical Tenderness noted in 55.9% patients and Forniceal tenderness in half of patients.

### Rupture status

2/3<sup>rd</sup> of patients presented with ruptured ectopic and hemoperitoneum. 1/3<sup>rd</sup> is unruptured ectopic.

### Patient management

Majority patients (91.3%) were managed surgically. Of these, Emergency laparotomy and salpingectomy in 71%, Salpingo-oophorectomy in 8.7% and laparoscopic salpingectomy in 11.6%. Three patients were managed conservatively. Blood transfusion was required in 21 patients. There were no deaths due to ectopic pregnancy in study period.

## Discussion

Incidence of ectopic pregnancy in our study is 0.87%, which is in accordance with previous studies in developing countries [2,5]. However incidence in the general population differs. In recent year's incidence have been increased with decrease in mortality due to ectopic pregnancy with advances in diagnosis and management procedures.

Abdominal pain was the predominant symptom followed by Amenorrhea and bleeding. Classical triad of symptoms was seen in 35.4% of patients. Other studies have reported it to be noted in 28%-95% of patients [7,8].

Table 1 Overall observation in section of patients.

Parameters	No. of cases	Percentage
<b>Age</b>		
15-20	4	11.8%
21-25	15	44.1%
26-30	9	26.5%
31-35	2	5.9%
36-40	4	11.8%
Total	34	100
<b>Parity</b>		
Nulli	9	26.5%
1	4	11.8%
2	9	26.5%
3	9	26.5%
4	3	8.8%
Total	34	100%
<b>Risk Factor</b>		
OCP	2	5.9%
Tubectomy	2	5.9%
Previous Abortion	6	17.7%
Previous Ectopic	2	5.9%
Infertility	2	5.9%
Previous C section	20	56%
<b>Symptoms (Normal)</b>		
Amenorrhea	26	76.5%
Pain abdomen	30	88.2%
Bleeding	20	58.8%
Others	9	26.5%
<b>General Physical Examination (Symptoms)</b>		
Pallor	26	76.5%
Shock	2	5.9%
None	8	23.5%
Fever	2	5.9%
<b>Abdominal Examination Findings</b>		
Tenderness	27	79.4%
Distension	5	14.7%
None	4	11.8%
Guarding	5	14.7%
<b>Per Speculum Examination</b>		
<b>Bleeding</b>		
Absent	14	41.2%
Present	20	58.5%
Total	34	100%
<b>Cervical Tenderness</b>		
Present	19	55.9%
Absent	15	44.1%
<b>Forniceal Tenderness</b>		
Absent	17	50%
Present	17	50%
<b>Rupture Status</b>		
Ruptured	23	67.6%
Unruptured	11	32.4%
<b>Diagnostic Studies</b>		
Laparotomy Unilateral Salpingectomy	24	71%

Parameters	No. of cases	Percentage
Salpingo-oophorectomy	3	8.7%
Laparoscopic Unilateral Salpingectomy	4	11.6%
Medical Management	3	8.7%
<b>Blood Transfusion</b>		
Done	21	61.8%
Not done	13	38.2%

Among the risk factors, Previous C section history is seen in more than half (56%) of the women. History of abortion was there in 6 patients (17.7%) of which 4 patients underwent dilatation and curettage. However studies from other regions have reported abortions as the major risk factor for EP in contrary to our study [5,9,10]. This can be explained by higher caesarian section rate in our state. There is History of tubectomy in two patients. Hence a pregnancy test should be performed in all suspected cases irrespective of tubal sterilization status [11]. Other risk factors were history of ectopic pregnancy and Infertility treatment (5.9% each).

Out of 11 unruptured ectopic pregnancies, 3 have been medically managed and other cases did not fit into the criteria for medical management. 2/3<sup>rd</sup> of patients (67.6%) presented with ruptured ectopic gestation. The higher rate can be explained by delayed presentation to our hospital as it is a referral center. Many previous studies reported similarly higher percentage of ruptured cases.

Most of the patients managed surgically (91.3%). Laparotomy and salpingectomy was the most commonly performed procedure. Laparoscopic salpingectomy (11.6%) was done in unruptured cases and few selected ruptured ectopic patients. A recent study revealed that there is no additional advantage with laparoscopic approach than laparotomy and is not a better option in terms of tubal patency and intrauterine pregnancy rates [12]. Most of the studies reported similarly higher rate of surgical management [2,5,13]. 21 women (61.8%) required blood transfusion which is

## References

- Singh S, Pukale RS, Mahendra G, Vijayalakshmi S (2014) A clinical study of ectopic pregnancy in a rural setup: A two year survey. *Natl J Med Res* 4: 37-39.
- Khaleeque F, Siddiqui RI, Jafarey SN (2001) Ectopic pregnancies: A three year study. *J Pak Med Assoc* 51: 240-243.
- Udigwe GO, Umeononihu OS, Mbachu II (2010) Ectopic pregnancy: A 5 year review of cases at Nnamdi Azikiwe University Teaching Hospital (NAUTH) Nnewi. *Nig Med J* 51: 160.
- Kirk E, Bottomley C, Bourne T (2014) Diagnosing ectopic pregnancy and current concepts in the management of pregnancy of unknown location. *Hum Reprod Update* 20: 250-261.
- Igwegbe A, Eleje G, Okpala B (2013) An appraisal of the management of ectopic pregnancy in a Nigerian tertiary hospital. *Ann Med Health Sci Res* 3: 166.

similar to that in other studies. Ultrasound being the mainstay of diagnosis, its availability at point of care allows early diagnosis and reduced morbidity [14].

There were no deaths reported during study period. Many studies reported a mortality rate of EP [2,5]. Mortality is mainly due to delayed diagnosis and hemoperitoneum. Recent reports affirm that early diagnosis is a key factor for effective non-surgical management. Horne et al. [15] suggested the use of serum Placental Growth Factor (PGF) to differentiate between intra uterine pregnancy and EP. They concluded that serum PGF was undetectable in women with tubal ectopic pregnancies compared to viable intra uterine pregnancies. Cabar et al. [16] correlated the levels of serum Vascular Endothelial Growth Factor (VEGF) and ultrasound findings in EP and concluded that serum VEGF was raised in EP with cardiac activity. Further studies are required to evaluate the clinical utility of these markers.

A recent study suggested use of letrozole (an aromatase inhibitor) in medical management of ectopic management, which is as effective as methotrexate (Chemotoxic agent). Metabolic and hormonal effects of these two medications are different; methotrexate elevates liver enzymes and decreases platelet count. Hormonal profile was more favorable in letrozole. HCG levels declines more rapidly with letrozole. After three months of medical treatment, letrozole does not affect the AMH levels where as methotrexate decreases the AMH levels affecting the fertility.

## Conclusion

Gynecologists should have comprehensive knowledge about the clinical presentation and risk factors of EP. If suspected clinically, encouraging women to undergo early ultrasonography allows early detection of EP and can be managed medically or fertility sparing surgical procedures.

## Limitations

Our study is limited by its retrospective nature and small sample size.

- Shah P, Shah S, Kutty RV, Modi D (2014) Changing epidemiology of maternal mortality in rural India: Time to reset strategies for MDG-5. *Trop Med Int Heal* 19: 568-575.
- Yadav K, Namdeo A, Bhargava M (2013) A retrospective and prospective study of maternal mortality in a rural tertiary care hospital of central India. *Ind J Comm Health* 25: 23-27.
- Jabbar FA, Al-Wakeel M (1980) A study of 45 cases of ectopic pregnancy. *Int J Gynecol Obstet* 18: 214-217.
- Shobeiri F, Tehranian N, Nazari M (2014) Trend of ectopic pregnancy and its main determinants in Hamadan province, Iran (2000-2010). *BMC Res Notes* 17: 7-13.
- Rather MS, Mufti S, Rangrez RA, Mufti KM, Rather SM (2012) Ectopic pregnancy: An analysis of 114 cases. *17: 21-24.*
- NFHS (2020) National Family Health Survey.
- Saranovic M, Vasiljevic M, Prorocic M, Macut ND, Filipovic T (2014).

- Ectopic pregnancy and laparoscopy. *Clin Exp Obstet Gynecol* 41: 276-279.
- 13 Cornelius AC, Onyegbule A, Onyema, Uchenna ET, Duke OA (2014) A five year review of ectopic pregnancy at Federal Medical Centre, Owerri, South East, Nigeria. *Niger J Med* 23: 207-212.
- 14 French S, Henry T, Williams EW (2014) Evaluation of waiting times and sonographic findings in patients with first trimester vaginal bleeding at the University Hospital of the West Indies: Can emergency department ultrasound make a difference? *West Indian Med J* 63: 247-251.
- 15 Horne AW, Shaw JLV, Murdoch A, McDonald SE, Williams AR, et al. (2011) Placental growth factor: A promising diagnostic biomarker for tubal ectopic pregnancy. *J Clin Endocrinol Metab* 96: E104-E108.
- 16 Cabar FR, Pereira PP, Schultz R, Francisco RP, Zugaib M (2015) Association between ultrasound findings and serum levels of vascular endothelial growth factor in ampullary pregnancy. *Fertil Steril* 103: 734-737.