

AGCHABADI DISTRICT OF THE NEW GARADOLAG VILLAGE ADMINISTRATIVE AREA EARTH COVER STUDY

PAE Z.H.Aliyev*

Department of Soil Science and Agrochemistry of ANAS, Somalia

*Corresponding author: PAE Z.H.Aliyev, Department of Soil Science and Agrochemistry of ANAS, Somalia. Tel: +252634818757; E-mail: Volqa_5@mail.ru

Received date: February 03, 2021; Accepted date: February 22, 2021; Published date: March 04, 2021

Citation: Aliyev PZH (2021) Agchabadi District of The New Garadolag Village Administrative Area Earth Cover Study, polisci, Vol. 7 No.1.

Copyright: © 2021 Aliyev PZH. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

It is known from the brief nature of the territory of the research district that the date of establishment of Agjabadi district - 08.08.1930 The territory - 1.76 thousand sq.m. km Population - 136.8 thousand people (January 1, 2020) Population density - 1 sq. km. km 78 people (January 1, 2020). Agjabedi district is located in Mil and Garabagh plains of Kur-Araz lowland. Agjabedi city and Sarvanlar, Kurds, Minakhorlu, Garavelli, Galabadin, Avshar, Mughanli, Shahsevan, Koyuk, Tazakend, Khojavend, Salmanbeyli, Ashagi Avshar, Hindarkh, Imamgulu-beyli, Mirzahagverdili, Sarijali, Balakhrizli, Taynaq, Poinag, Sharafkhanli, Shotlanli, Shenlik, Husulu, Ranjbarlar, Boyat, Hajilar, Garakhanli, Hajibadalli, Pariogullar, Agabeyli, Garadolag, Mehrabli, Kabirli, Aran, Yeni Garadolag, Najaf-gulubeyli, Qiyameddinli, Shahsevan-Tazakand, Shahsevan-Tazakand, Includes Jafarbeyli and Yukhari Qiyameddinli villages. The relief of the region is flat, gradually rising from the north-east to the south-west. The surface of the area is composed of continental-alluvial and marine sediments of the Anthropogenic system. There is a clay deposit. The climate is temperate hot, dry subtropical. The average temperature is 1.2-1.7 ° C in January and 25-26 ° C in July. Annual precipitation is 300-500 mm. The river network is sparse. The Kura River flows along the north-eastern border, and the Gargar River flows through the central part. The Upper Karabakh canal passes through the region. There is a salt lake in the area. Gray-meadow, gray, meadow-gray soils are widespread. In the central part, saline and saline soils are found. The plants are of steppe and semi-desert type. There are bushes and sparse Tugay forests on the banks of the Kura River. Animals: gazelle, wolf, wild boar, jackal, fox, swamp beaver, badger, gray rabbit, Asia Minor sand mouse, etc. Birds: turkey, pigeon, black grouse, pheasant, etc. Aggol National Park is located in Agjabadi district. 15 species of animals, 20 species of fish and 40 species of plants have been recorded here. Agjabadi is mainly an agricultural region. Fruit growing, cotton growing, grain growing, silkworm breeding, animal husbandry, etc. **developed.**

Keywords: soil formation, soil cover, gray-meadow, light gray-meadow, heavy clayey, light clayey salinity, salinity, erosion, etc.

Introduction

The establishment of Garadolag village is connected with the name of Garadolag tribe. In the 18th century, Panahali khan moved some of the Garadolags to Karabakh as part of the Kangarli [1]. As a result of the Kura flood in 1938, the Garadolags were relocated from the Kurkiragi areas where they settled to the area where they now live. Here they created Garadolag and then Yeni Garadolag villages. Garadolag village has an area of 32.7 square kilometers and a population of 3139 people [2]. The main occupations of the population are agriculture and animal husbandry [3].

The course of the study:

In accordance with the requirements of the existing agro-industrial grouping, in April 2017, a soil survey was conducted in the administrative territory of Yeni Garadolag village of Agjabedi district and the following was determined:

The total area of the surveyed area is 2180.77 ha. Land survey works covered an area of 1802.75 hectares [4].

The area is divided into the following natural areas:

Planting 1760.78 ha

Dinc 15.53 ha

Clean pasture 26.00 ha

Perennial plantings 0.44

ha Other lands 378.02 ha

During the study, 82 sections were excavated in the area and morphological features were described in genetic layers [5]. Soil samples were taken from the excavated sections and the following analytical-laboratory researches were carried out on them and the results were analyzed:

1. Hygroscopic moisture - by thermal method
2. Granulometric composition - by Kaczynski's pipette method
3. General humus - by the method of Tyurin
4. Total nitrogen - By calculation
5. Carbonate - With a calcimeter device

6. Absorbed Ca and Mg - by Ivanov method

7. Absorbed Na - by Hedroysts method

8. pH water suspension - with pH meter

Thus, based on the results of field soil research and laboratory analysis, a soil map was prepared on a topographic basis and a report was written.

Archival materials were used in compiling the maps and writing the report.

From the scale of Professor RH Mammadov in determining the granulometric composition was used.

Natural conditions

Geographical position

The studied area is bounded on the north-west and north by Avshar village, on the east by Garadolag village, on the south-east by Mehrabli village, on the south by Kabirli village, on the south-west by Aran village and on the west. Avshar and Aran villages of winter pasture No. 104 are bordered by territorial lands.

Relief

Relief, as a structure of the earth's surface, is directly involved in the formation of land cover as a factor in soil formation. It plays an important role in changing chemical and biological processes, hydrothermal regime and microclimate [6]. Thus, the distribution of solar energy and atmospheric sediments is directly related to relief. The relief of the studied area consists of sloping and wavy plains.

Climate

Climate is one of the important factors as a factor in soil formation. Agjabedi district is located in the Kur-Araz lowland, in the western part of the Mil plain. The climate of the area is temperate hot semi-desert and dry steppes with dry summers, weak humidification is characterized by hot summers. The average temperature in January is 1.8 0C, the average temperature in July is 260C [7]. The average annual relative humidity is 73%. The annual rainfall is 332 mm, mainly in spring and autumn. 980 mm of possible evaporation from the surface cover per year. The average annual soil surface temperature is 180C.

Vegetation

Vegetation is a key factor in the process of soil formation and the formation of soil cover. The increase in soil fertility with the formation of organic matter depends on the density of vegetation.

Maintaining normal soil moisture, reducing the washing effect of water, preventing the formation and development of the erosion process are closely related to vegetation.

In the area we studied, we found sagebrush, wild clover, cattail, solid guramid, yellow flower, carnation, reed, yaghtikan, invitation, meadow, etc. plants are widespread.

Cultivated crops are also grown in the area.

Soil-forming rocks

Soil-forming rocks affect the granulometric composition, chemical and mineralogical composition of the soil, causing the formation of soil profile and genetic layers [8]. The chemical composition of the parent rock plays an important role in the process of soil formation. The richer the parent rock, the better the quality of the soil formed on it.

Thus, the soils of the area under study were formed on alluvial sediments.



Figure 1: Description of the land plot in the area

Ground cover

According to the results of field soil research and laboratory analysis, the following soil types and subtypes are widespread in the area.

- I. Gray-meadow soils
- II. Light gray-meadow soils

I. Gray-meadow soils

Gray-meadow soils cover 761.47 ha or 34.92% of the total area, spreading to the west and east of the study area.

These soils are due to their granulometric composition and soil layer thickness; 1) Divided into heavy clayey, thick, gray- meadow species.

To get acquainted with the characteristic morphological features of these soils, we give a description of section 29 excavated in the area.

0-23 cm gray, topavari, heavy clayey, solid, tubers and rhizomes, boiling, dry, the transition is 23-54 cm light gray, topavari, light clay, less solid, root residue, boils, slightly moist, the transition is gradual 54-89 cm light gray, small clump, light clay, less solid, insect tracts, boils, less moisture, the transition is 89-127 cm grayish, small topavari, light clay, soft, rust spots, boils, less moisture, the transition is gradual 127-156 cm straw, not selected, light clay, soft, carbonate stains, boiling, wet [9]. It is clear from the morphological description of the section that the color of these soils is gray in the upper layer, light gray and gray in the middle layer, and straw in the lower layer. The structure is not selected on the top layer, topavari and small topavari on the middle layer, and on the bottom layer. The granulometric composition was heavy clayey, and the length of the profile was wool the rose is clay. The density is hard on the top layer and less hard and soft on the bottom layers.

Roots and rhizomes, root residues, insect tracts, rust and carbonate stains are found along the profile from new derivatives and marshes.

These soils are boiled under the influence of 10% hydrochloric acid (HCl).

Humidity is dry in the upper layer, slightly moist in the middle layers, and moist in the lower layer. Transitions to genetic layers are clear and gradual throughout the profile.

According to the results of laboratory analysis, the granulometric composition of gray-meadow soils is heavily clayey. Thus, the amount of physical clay in these soils is 47.80-49.18% in the upper layers, and the profile length is 47.80-55.84%

The hygroscopic humidity of the main components varies between 4.1-5.3% throughout the profile.

Total humus is 2.07-2.23% in the upper layers, and the profile length is 1.09-2.23%. According to the total humus, the total nitrogen content is 0.10-0.17%. The amount of carbonate is 10.69-13.68% throughout the profile. The pH in the water suspension is 8.20-8.33 units, which indicates that these soils are alkaline.

The total amount of absorbed bases in gray-meadow soils is 28.40-32.95 mg. As a percentage, the Ca base is 62.08-63.74%, the Mg cation is 33.33-35.24%, and the Na cation is 2.68-3.17%.

II. Light gray-meadow soils

Light gray-meadow soils cover 1041.28 ha or 47.75% of the total area, spreading to the north and south of the study area.

These soils are divided into the following types according to their granulometric composition and thickness of the soil layer.

- 2) Light clay, thick, light gray-meadow
- 3) Heavy clayey, thick, light gray-meadow

To get acquainted with the characteristic morphological features of these soils, we give a description of section 55 excavated in the area.

0-22 cm light gray, topavari, light clay, solid, root and roots, boiling, dry, clear transition.

22-55 cm light gray, topavari, light clayey, less solid, root residue, boil, dry u, the transition is gradual.

55-88 cm grayish, small clumpy, lightly clayey, less solid, vertical cracks, boils, less moisture, gradual transition.

88-122 cm grayish, small topavari, light clay, soft, rust stains, boils, less moisture, gradual transition.

122-155 cm straw, not selected, light clay, soft, carbonate spots, boils, less moisture.

It is clear from the morphological description of the section that the color of these soils is light gray in the upper layer, light gray and gray in the middle layers, and straw in the lower layer. The structure is not selected on the top layer, topavari and small topavari on the middle layer, and on the bottom layer. The granulometric composition is light clay throughout the profile. The density is hard on the top layer and less hard and soft on the bottom layers. Roots and rhizomes, root residue, vertical cracks, rust and carbonate stains are found along the profile from new derivatives and alloys. These soils are boiled under the influence of 10% hydrochloric acid (HCl). Humidity is dry in the upper layers and less moist in the lower layers. Transitions to genetic layers are clear and gradual throughout the profile.

According to the results of laboratory analysis, the granulometric composition of light gray-meadow soils is light clayey and heavy clayey. Thus, the amount of physical clay in these soils is 45.16-52.08% in the upper layers, and the profile length is 45.16-55.96%.

Conclusion

The study revealed that the presence of many structural aggregates larger than 1 mm in the soil, its physical, physico-chemical, etc. indicators show that they are resistant to erosion. Decreased water-resistant structural aggregates in eroded soils, easy dusting of the soil, compaction, destruction of non-capillary pores, increased surface water flow, making it difficult for atmospheric sediments to absorb into the soil. As a result, the amount of washed soil throughout the area has increased. For this reason, due to the loss of large amounts of nutrients along with the washed soils, the concentration of microelements in the soil solution decreased and the transpiration of water by plants increased. Depending on the shape of the terrain, most of the atmospheric water in eroded soils flows along the slope, loses nutrients, and the water absorbed into the soil is used for physical evaporation and transpiration by plants. It can be concluded that the development and intensity of erosion can vary in one direction or another, depending on the granulometric composition, which is closely related to its granulometric composition, which leads to a lack of oxygen in the soil pores and difficulty in respiration of plant roots. The role of granulometric composition in changing the rate of erosion on

the soil surface and its washing intensity is great. Studies have shown that heavy granulometric soils with high cohesiveness and poor water permeability are more resistant to erosion than light granulometric soils, which are less resistant to heavy rains and even on very sloping slopes and have good water permeability. The light granulometric soils present in the less inclined parts of the slope absorb water well in light rains, which reduces surface water flow and soil leaching.

PRACTICAL PROPOSALS FOR ECONOMY

Object of research

Given the fact that the existing soils in the zone are mainly weakly and moderately, and partially severely eroded under the planting and grazing areas, the complex implementation of the following proposals is necessary:

1. All agro-technical measures should be carried out in the direction of the width of the slope by organizing grassland rotation in the sown areas, the areas should be cleared of stones, organic and mineral fertilizers should be applied in time, taking into account the degree of erosion. Slopes up to 11° should be plowed directly in the direction of the slope width. Slopes 11°-19° should be plowed by semi-tillage, slopes 19°-27° should be plowed in strips (8-10 m plow, 3-5 m grass strip), slopes more than 27° should be used only under fruit trees. Trees should be planted in the direction of the width of the slope by means of pits and ditches.

2. Deep loosening of strips in arable lands improves the absorption of atmospheric water by the soil, weakens the surface water flow, increases soil moisture reserves, reduces erosion and significantly increases the productivity of agricultural crops.

3. In moderately eroded fields, perennial grasses should be planted for 3-5 years, high doses of organic and mineral fertilizers should be applied, plantation plowing should be carried out in the third and fifth years without harvesting, and then used under cereals and vegetables.

4. In order to make efficient use of pastures, they should be divided into at least 5-6 equal parts and the cattle should move to the next part after grazing in each part for 4-5 days. By the time the last part is reached, the grass plants in the first part have grown enough and are in a condition suitable for grazing. Thus, the pasture rotation is organized.

5. Slightly eroded pastures should be cleared of stones, sparse areas should be sown with grass seeds in accordance with local conditions, organic and mineral fertilizers should be applied, alternating grazing should be applied, 1 head of cattle or 7-8 heads of small cattle should be grazed per hectare.

6. In moderately eroded pastures, the grazing rate should be reduced by 50% (0.5 head of cattle, 3.5-4.0 head of small cattle should be grazed on 1 ha of pasture), grazing should be stopped on rainy days, grass seeds should be sown in areas with sparse grass cover, organic and mineral fertilizers should be applied.

7. Rehabilitation methods, especially the "corridor" method, are more appropriate to rehabilitate low-value forests and shrubs. When applying this method, care should be taken to ensure that the soil preparation is in the form of plots.

References

1. Aliyev GA. (1978) Soils of the big Caucasus within the Azerbaijan SSR, Baku, Elm ", 157.
2. Aliyev BH, Aliyev ZH, Aliyev IN. Problems of erosion in Azerbaijan and Said Russian ways of its solution.
3. Aliev ZH. (1999) Technique and technology of low-intensity irrigation in the conditions of the mountainous region of Azerbaijan Elm Publishing House Baku. 220.
4. Aliev ZH. (2005) Irrigated agriculture in the mountainous and foothill regions of Azerbaijan. Publishing House "Ziya-Nurlan". 330.
5. Aliyev BH, Aliev IN. (2004) Some problems of agriculture in Azerbaijan and ways to solve them Ziya-Nurlan Publishing House. 572.
6. Araev HA. (1998) Content of micronutrients in the main soil-forming rocks of the Lesser Caucasus of Azerbaijan. DAN Azerbaijan, Ed. Elm, Baku, 59-63.
7. Qiyasi HA. (2000) Effect of erosion on soil fertility. Works of the Institute of Erosion and Irrigation. Volume II. Nurlan Publishing House. Baku, 89-102.
8. Glushko EV, Ermakov YG, Serebrov AA. (1986) Application of space images in the study of the development of modern landscapes. Exploration of the Earth from space. Izd. Science, Moscow, № 3, 21-28.
9. Mustafayev XM. (1978) Measures to combat water and wind erosion. Baku, 228.