

Correlation Between Psychological Distress and Internet Addiction Among Adolescents During Covid-19 Period

Abstract

Background: The virtual environment, facilitated through the use of the Internet, is a well acknowledged socio-economic-cultural reality of the present times, especially in the wake of COVID-19 related precautionary measures and restriction related to real time interactions. At the same time, poor psychological health can be a trigger for using internet as a means of escape from boredom, loneliness, anxiety, sadness, and frustration, a trend which seems to be rising during the current COVID-19 period. Long hours of internet use, with no outdoors do have consequences for the psychological health of an adolescent.

Objective: The present study explores the correlation between psychological distress and internet addiction among adolescents.

Materials and Methods: This descriptive and observational study was conducted in the Department of Psychiatry, National Institute of Mental Health & Hospital, Dhaka, Bangladesh from January 2020 December 2020. In survey, a sample of 51 adolescents in the age range of 13 to 18 years was administered Internet Addiction Test. The correlation between Internet Addiction and Psychological Distress, as measured through the levels of anxiety, depression and loss of behavioral/emotional control, was examined using Pearson Product Moment Correlation Method.

Results: The total number of uncontrolled users of internet (moderately and severely addicted combined) constituted about two thirds (66.7%) of the sample population. Moderate to severe Psychological Distress was discovered in 71.7% of the sample population. There were no gender differences in Internet Addiction. Internet Addiction positively correlated with Psychological Distress ($r=0.82$; $p<0.01$).

Conclusion: The assessment of Internet usage can be helpful in the treatment of psychological problems among adolescents. In a similar vein, for the treatment of Internet addicted adolescents, the evaluation of psychological distress, particularly depression and anxiety, must be considered. Effective counseling and intervention programs need to be designed and implemented for teaching healthy, moderate and non-addictive internet use for improving the overall psychological well-being of the adolescents.

Keywords: Adolescents; Depression; COVID-19; Psychological Distress

Received: July 06, 2021, **Accepted:** July 08, 2021, **Published:** augst 20, 2021

Introduction

The virtual environment, facilitated through the use of the Internet, is a well acknowledged socio-economic-cultural reality of the present times, especially in the wake of COVID-19 related precautionary measures and restriction related to real time interactions. In the face of social isolation and online schooling,

there is an exponential rise in the screen time and internet usage by the young generation. The Data from the 2004 National Health Information Survey (2004) found that over 1 in 10 (11.6%) adolescents aged 12-17 years had serious behavioral or mental health difficulties [1]. Found that the levels of depression and suicide ideation were highest in the Internet-addicts group of Korean adolescents, the prevalence of internet addiction not

**Niaz Mohammad Khan^{1*},
MSI Mullick², Sharmin
Hussain³ and Uddin MMJ**

1 Associate Professor of Psychiatry,
National Institute of Mental Health,
Dhaka, Bangladesh

2 Professor of Child and Adolescent
Psychiatry, Bangabandhu Sheikh Mujib
Medical University, Dhaka, Bangladesh

3 Junior Consultant, Pediatrics, and FCPS
Trainee of Pediatric Neurology, National
Institute of Neurosciences (NINS), Shere
Bangla Nagar, Dhaka, Bangladesh

4 Associate Professor (Psychiatry), National
Institute of Nuro Sciences (NINS), Dhaka,
Bangladesh

*Corresponding author:

Niaz Mohammad Khan

 nmadil27@grnail.com

Department Psychiatry, National Institute of
Mental Health, Dhaka, Bangladesh

Citation: Khan NM, Mullick MSI, Hussain
S, Uddin MMJ (2021) Correlation Between
Psychological Distress and Internet
Addiction Among Adolescents During
Covid-19 Period. Br J Res Vol. 8 No. 8: 39.

varying with gender [2]. In a two-wave panel household survey, found a positive correlation found between number of symptoms of internet addiction and depressive symptoms among a sample of 208 adolescents, aged 15-19 years, in their review of the research published in Medline and Pub Med from 2000-2009, discovered a high co-morbidity of Internet addiction with affective disorders and anxiety disorders [3]. Found that suicidal behaviors, depression, anxiety, conduct problems and hyperactivity/inattention were significant and independent predictors of pathological internet use [4]. The general prevailing atmosphere of isolation, anxiety, and fear is leading to addictive behaviors like compulsive eating, substance use, and compulsive internet use, as a means of distraction from the uncomfortable feelings arising in the current situation. There is no doubt that the adolescents today are gravely affected by excessive internet use, by what develops into an unhealthy and uncontrollable attachment to internet use. At the same time, addiction to the Internet and electronic devices might just be the tip of an ice-berg, a larger psychological problem underlying the whole phenomenon of Internet Addiction. Internet addiction, compulsive internet use, problematic internet use and pathological internet use, are some of the concerns that have merited the attention of clinical psychologists and mental health professionals all over the world. This compulsive and addictive use of the internet can have significant influence on the psychological health of the emotionally most vulnerable individuals of the society, that is, the adolescents. Therefore, research on the different aspects of psychological health of adolescents in relation to internet addiction is one of the most pressing needs in the field of research in community mental health and psychopathology. This need becomes even more profound now than ever because of the escalation in digital consumption due to COVID-19 related stay-at-home mandates and the associated psychological concerns arising due to anxiety, insecurity, frustration, deprivation, and boredom. The term internet addiction was proposed by Dr. Ivan Goldberg in 1995 for a maladaptive pattern of Internet use, leading to clinically significant impairment or distress [5]. The addiction is characterized by the use of computers that takes up so much time as to cause discomfort or decreased occupational, academic, social, work-related, family-related, financial, psychological, or physiological functioning [6]. According to Dr. Young (1996), Internet addiction is any online-related, compulsive behavior which interferes with normal living and causes severe stress on family, friends, loved ones, and one's work environment [7]. Young (1999) is of the opinion that Internet addiction could actually result from another existing disorder like depression, stress or anxiety [7]. The state of emotional suffering characterized by symptoms of depression, e.g., lost interest, sadness, hopelessness; and anxiety, e.g., restlessness, feeling tense; has been defined as psychological distress [8]. Psychological distress is widely used as an indicator of the mental health of the population in public health, in population surveys and in epidemiological studies and, as an outcome, in clinical trials and intervention studies [9]. Veit and Ware (1983) conceptualized psychological distress as comprising of anxiety, depression and loss of behavioral/emotional control [10]. Parent's Guide to Teen Depression opines that teens may go online to escape their problems, but excessive computer use only increases their isolation, making them more depressed.

Bajaj & Sheoran (2016) stated that this addiction to internet can be viewed as a double edged sword [11]. It can be conceived to be an outcome of psychological distress among adolescents, as most of the addictions are. Moreover, the addiction to internet itself may have severe implications for the mental health of the young adolescents [11]. Internet addiction was associated with gender as well as with anxiety, depression, and loss of emotional/behavioral control. Sharma & Sharma (2018) found that the Psychological Well-Being of college students was negatively affected by internet addiction [12]. Based on a study involving 1086 engineering students, reported that 27.1% of the participants met criterion for mild addictive internet use, 9.7% for moderate addictive internet use, and 0.4% for severe addiction to internet [13]. The researchers also reported that psychological distress (depressive symptoms) predicted Internet Addiction. In a comparative study, discovered that the COVID-19 outbreak has had a significant effect on adolescent internet use and psychosocial well-being [14]. According to (2020), the frequency and duration of recreational internet use, and the frequency of stay-up internet use internet usage has grown during the COVID-19 epidemic.

Material and Methods

This descriptive and observational study was conducted in the Department of Psychiatry, National Institute of Mental Health & Hospital, Dhaka, Bangladesh from January 2020 December 2020. In survey, a sample of 51 adolescents in the age range of 13 to 18 years was administered Internet Addiction Test. The correlation between Internet Addiction and Psychological Distress, as measured through the levels of anxiety, depression and loss of behavioral/emotional control, was examined using Pearson Product Moment Correlation Method. The study uses a descriptive survey method. The participants consist of a total of 51 adolescents (Mean Age=15.57, SD= +1.42), selected through quota sampling from different city of Bangladesh. Internet Addiction Test (IAT) is a 20-item questionnaire, answered in a five-point Likert scale and covers six factors, that is, Salience, Excess Use, Neglecting Work, Anticipation, Lack of Self-control and Neglecting Social Life. Developed by Dr. Kimberly Young, the IAT measures; mild, moderate and severe levels of internet addiction. Individuals who score 0-30 are classified as average online user, or normal Internet user; 31-49 as controlled users, or mild Internet Addiction; 50-79 as experiencing occasional or frequent problems because of uncontrolled use of the Internet, or moderate Internet Addiction; and 80-100 as having significant problems in life because of the uncontrolled use of the Internet, or Severe Internet Addiction [15]. The Pearson's r for the correlations among the six factors ranges from r=0.62 to r=0.226 [16]. These factors show good to moderate internal consistency (alpha coefficients 0.54 to 0.82).

Mental Health Inventory (MHI-38)

Psychological Distress was measured through the Psychological Distress Subscale of Mental Health Inventory (MHI-38). The MHI has 38 items scored on a six-point scale (range 1-6) except two items 9 and 28 which are scored on a five-point scale (range 1-5). The inventory measures two global Scales-Psychological

Distress and Psychological Well-being; as well as global Mental Health Index. For the purpose of this study, the Psychological Distress (PD) subscale was used. There are three dimensions of PD- Anxiety, Depression, and Loss of Behavioral/Emotional Control. The Cronbach alpha for Psychological Distress has been estimated at 0.94 [10].

Procedure

A total of 51 adolescents (26 males and 25 females) in the age group of 13-18 years were administered the Internet Addiction Test and Psychological Distress Subscale of Mental Health Inventory through Google forms. The statistical techniques of Mean, Standard Deviation, Percentage and Product Moment Correlation were used to analyze and interpret the data.

Results

As seen in the Mean and SD of Internet Addiction among adolescents were found to be 57.09 and 20.68 respectively (**Table 1**). **Table 2** shows that out of the 51 adolescents that constituted the sample population, only 10 (19.60%) were found to be using the Internet within normal limits. While 15 adolescents (29.47%) were mildly addicted to the Internet, 17 adolescents (33.3%) met the criteria for moderate addiction, and 8 adolescents (15.69%) were severely addicted. **Table 3** summarizes the differences in the Mean IAT scores of the male and female adolescents, the mean scores of female adolescents being slightly higher at 58.45 than

the mean of male adolescents at 55.73. The t-value is 1.14 which is not significant. Hence, our null hypothesis that there exists no significant difference in Internet Addiction among the male and female adolescents is not rejected.

Table 4 summarizes the Mean Scores of Psychological Distress (PD) variables among adolescents. The mean score on Anxiety was 29.44, while the numbers of adolescents in normal, moderate and high range of anxiety were 21%, 39%, and 39% respectively. On Depression, the mean score was 12.89. Approximately one third of the populations (39%) were in normal range, while 39% were moderately depressed, and 21% met the criteria for severely depressed. Loss of Behavioral/Emotional Control showed a mean of 27.08, with the normal, moderate, and severe range percentages being 21%, 39%, and 39% respectively. The Psychological Distress mean was found to be 75.27. It was found to be within acceptable range in 19% of the adolescents, while 39% showed moderate levels, and 41% showed severe levels of Psychological Distress. Data presented in **Table 5** shows that the mean PD scores of female adolescents were higher than that of the male adolescents, the scores being 77.21 and 73.33 respectively. The t-value for this difference was found to be 1.86 which is not significant. Hence, our null hypothesis that there exists no significant difference in Psychological Distress among the male and female adolescents is not rejected.

Data presented in **Table 6** indicate that the value of Pearson Product Moment Correlation Coefficient between Anxiety and

Table 1: Mean and SD of Internet Addiction Test (IAT) Scores among Adolescents.

Variable	Number of adolescents	Mean age (in years)	Mean iat score	SD
Internet addiction	51	15.57	57.09	20.68

Table 2: Number and Percentage of Adolescents Showing Different Levels of Internet Addiction.

Levels of internet addiction	Score Range	Number of Adolescents	Percentage
Normal (average users)	0-30	10	19.60
Mild addiction (controlled users)	31-49	15	29.41
Moderate addiction (uncontrolled users with occasional or frequent problems)	50-79	17	33.33
Severe addiction (uncontrolled users with significant problems)	80-100	8	15.69
Total	0-100	51	100

Table 3: Significance of Difference between IAT Means of Male and Female Adolescents

Gender	Mean age	Mean iat scores	sd	sed	t'-value	Remark
Male adolescents (n=26)	15.52	55.73	19.96	2.39	1.14	p>0.01

Table values of t' (df=298) at 0.05 and 0.01 level of confidence are 1.97 and 2.59 respectively.

Table 4: Mean, SD and Percentage of Psychological Distress (PD) Variables among Adolescents.

Psychological distress variables	Score range	Sample mean	Sd	Levels of Psychological distress	Number of adolescents	Percentage
Anxiety (A)	9 to 54	29.44	6.81	Normal	11	21.56
				Moderate	20	39.21
				High	20	39.21
Depression (D)	4 to 23	12.89	4.34	Normal	20	39.21
				Moderate	20	39.21
				High	11	21.56
Loss of behavioral/emotional control (B)	9 to 53	27.08	7.13	Normal	11	21.56
				Moderate	20	39.21
				High	20	39.21
Psychological distress (PD)	24 to 142	75.27	18.07	Acceptable	10	19.60
				Moderate	20	39.21
				High	21	41.17

Table 5: Significance of Difference between Psychological Distress (PD) Means of Male and Female Adolescents.

Gender	Mean age	Mean pd scores	SD	SED	t' -Value	Remark
Male adolescents (n=26)	15.52	73.33	17.89	2.08	1.86	P > 0.01
Female adolescents (n=25)	15.51	77.21	18.10			

Table values of t' (df=298) at 0.05 and 0.01 level of confidence are 1.97 and 2.59 respectively.

Table 6: Correlation between Internet Addiction and Psychological Distress among Adolescents.

Correlation variables		Mean	Sd	Correlation
Correlation between anxiety and internet addiction	Anxiety	29.44	6.81	0.73**
	Internet Addiction	57.09	20.68	
Correlation between depression and internet addiction	Depression	12.89	4.34	0.77**
	Internet Addiction	57.09	20.68	
Correlation between loss of Emotional/ behavioral control and internet addiction	Loss of Behavioral/ Emotional Control	27.08	7.13	0.78**
Correlation between psychological distress and internet addiction	Internet Addiction	57.09	20.68	
	Psychological Distress	75.27	18.07	0.82**

* Significant at 0.05 level

**Significant at 0.01 level

Internet Addiction was 0.73, between Depression and Internet Addiction was 0.77, between Losses of Behavioral/Emotional Control was 0.78. The Pearson Product Moment Correlation between Psychological Distress and Internet Addiction among adolescents was 0.82. These correlations are positive and are significant at 0.01 level of confidence.

Discussion

The results of the study revealed that 19.6% of the sampled population was making normal use of the Internet, 19.7% were mildly addicted and 29% were moderately addicted and another 33% were in Severe Addiction category. In a two-wave panel household survey, found a positive correlation found between number of symptoms of internet addiction and depressive symptoms among a sample of 208 adolescents, aged 15-19 years. In their review of the research published in Medline and Pub Med from 2000-2009, discovered a high co-morbidity of Internet addiction with affective disorders and anxiety disorders. Found that suicidal behaviors, depression, anxiety, conduct problems and hyperactivity/inattention were significant and independent predictors of pathological internet use. A descriptive, cross-sectional, correlational study by established the prevalence of Internet Addiction at 40% among 587 university students in Jordan [17]. Al-Gamel and colleagues also reported that Internet Addiction was associated with high mental distress among the students [17]. It revealed that Internet addiction mediated the association between social anxiety and poor psychosocial well-being among adolescents in China, Hong Kong, and Malaysia [18]. Kawa & Shafi discovered that male university students experienced more internet addiction and psychological distress as compared to the female university students and internet addiction was positively correlated with psychological distress [19]. In a study involving university undergraduate students, Uddin observed that 47.7% male and 44.5% female students showed severe Internet addiction, while 27.1% male and 33.9% female students showed moderate Internet addiction [20]. In a cross-sectional study conducted among 846 students of various faculties from Deemed University, Gedam reported that the total prevalence of internet addiction was 19.85%, with moderate

and severe addiction being 19.5% and 0.4%, respectively [21]. Internet addiction was associated with gender as well as with anxiety, depression, and loss of emotional/behavioral control. Sharma & Sharma (2018) found that the Psychological Well-Being of college students was negatively affected by internet addiction [12]. Based on a study involving 1086 engineering students, Anand reported that 27.1% of the participants met criterion for mild addictive internet use, 9.7% for moderate addictive internet use, and 0.4% for severe addiction to internet. In fact, the situation is quite alarming with the total number of uncontrolled users of internet (moderately and severely addicted combined) constituting about two thirds (66.7%) of the sample population. There were, however, no significant gender differences in Internet Addiction among the male and female adolescents. The present findings are in line with Kim who reported that the prevalence of Internet Addiction did not vary with gender. However, the results refute the findings of Kawa and Shafi and Gedam who had reported that Internet Addiction was associated with gender. It may be said that the trends are repeating, and pathological dependence on Internet use is increasing among the adolescents, irrespective of gender. Both the male and female populations are equally vulnerable. In case of Psychological Distress, more than one third of the sample population (35.3%), was in moderate range and another 36.3% population was found to be in the severe range. The female adolescents seem to be at a higher risk of Psychological Distress as compared to male adolescents though the differences with respect to gender are not statistically significant. The current study established significant positive correlation between Psychological Distress and Internet Addiction among adolescents ($r=0.82$). The findings are in line with the results of Gedam who confirmed that Internet addiction was associated with anxiety, depression, and loss of emotional/behavioral control. The present results also corroborate the findings Kim who reported that psychological distress (depressive symptoms) predicted Internet Addiction. Given the definite, high, significant positive correlation between Psychological Distress and Internet Addiction among adolescents, it is recommended that internet addiction should be viewed in relation with the psychological health of the adolescents. Adolescents screened for internet addiction should also be assessed for psychological

distress variables like anxiety, depression, and loss of behavioral/emotional control. A comprehensive psychological health program must involve appropriate interventions for psychological distress along with steps for internet de-addiction. Educational institutions and community mental health bodies should work in tandem to raise awareness regarding the healthy, balanced and non-addictive use of internet. Non-digital leisure and entertainment activities like reading, painting, sports, music, dance, etc. should be encouraged. Steps to develop strategies for prevention of internet addiction can go a long way in promoting the psychological well-being of the adolescents.

References

1. Parker J D, Akinbami L J, Woodruff T J (2009) Air pollution and childhood respiratory allergies in the United States. *Environ Health Perspect*, 117: 140-147.
2. Sheoran P (2021) Prevalence of Psychological Distress among Adolescents in relation to Internet Addiction during COVID-19 Times. *Acade Accel worl res* 8: 11-15.
3. Sharma A, Sharma R (2018) Internet addiction and psychological well-being among college students: A cross-sectional study from Central India. *J Family Med Prim Care* 7: 147.
4. Anand N, Jain P A, Prabhu S, Thomas C, Bhat A, et al. (2018) Internet use patterns, internet addiction, and psychological distress among engineering university students: A study from India. *Indian J Psychol Med*, 40: 458-467.
5. Fernandes B, Biswas U N, Mansukhani R T, Casarín A V (2020) The impact of COVID-19 lockdown on internet use and escapism in adolescents. *Revista de psicología clínica con niños y adolescentes* 7: 59-65.
6. Young K S, Yue X D, Ying L (2011). Prevalence estimates and etiologic models of Internet addiction. *Inter addi: A handb gui evalua treatm*, 3-17.
7. Sharma A, Sahu R, Kasar P K, Sharma R (2014) Internet addiction among professional courses students: A study from central India. *Int J Med Sci Public Health*, 3: 1069-1073.
8. Al-Gamal E, Alzayyat A, Ahmad M M (2016) Prevalence of internet addiction and its association with psychological distress and coping strategies among university students in Jordan. *Perspect Psychiatr Care*, 52: 49-61.
9. Lai C M, Mak K K, Watanabe H, Jeong J, Kim D, et al. (2015) The mediating role of Internet addiction in depression, social anxiety, and psychosocial well-being among adolescents in six Asian countries: a structural equation modelling approach. *Public health*, 129: 1224-1236.
10. Kawa M H, Shafi H (2015) Evaluation of internet addiction, impulsivity and psychological distress among university students. *Int J Clin Ther Diagn*, 3: 70-76.
11. Uddin M S, Al Mamun A, Iqbal M A, Nasrullah M, Asaduzzaman M, et al. (2016) Internet addiction disorder and its pathogenicity to psychological distress and depression among university students: A cross-sectional pilot study in Bangladesh. *Psychology* 7: 1126.
12. Gedam S R, Ghosh S, Modi L, Goyal A, Mansharamani H (2017) Study of internet addiction: Prevalence, pattern, and psychopathology among health professional undergraduates. *Indian J Soc Psychiatry*, 33: 305.
13. Kim K, Ryu E, Chon M Y, Yeun E J, Choi S Y, et al. (2006) Internet addiction in Korean adolescents and its relation to depression and suicidal ideation: a questionnaire survey. *Int J Nurs Stud* 43: 185-192.
14. Fu K W, Chan W S, Wong P W, Yip P S (2010) Internett-avhengighet: utbredelse, diskriminerende gyldighet og korrelasjoner blant ungdom i Hong Kong. *Br J Psykiatri* 196: 486-492.
15. Weinstein A, Lejoyeux M (2010) Internett-avhengighet eller overdreven bruk av Internett. *Am J Drug Alcohol Abuse* 36: 277-283.
16. Kaess M, Durkee T, Brunner R, Carli V, Parzer P, et al. (2014) Pathological Internet use among European adolescents: psychopathology and self-destructive behaviours. *Eur Child Adolesc Psychiatry* 23: 1093-1102.
17. Goel D, Subramanyam A, Kamath R (2013) A study on the prevalence of internet addiction and its association with psychopathology in Indian adolescents. *Indian J Psychiatry*, 55: 140.
18. Young K S, De Abreu C N (2010) Internet addiction: A handbook and guide to evaluation and treatment. John Wiley & Sons.
19. Mirowsky J, Ross C E (2002) Measurement for a human science. *J Health Soc Behav* 152-170.
20. LAbate L (Ed) (2012) Mental illnesses: understanding, prediction and control. BoD-Books on Demand.
21. Veit C T, Ware J E (1983) The structure of psychological distress and well-being in general populations. *J Consult Clin Psychol* 51: 730.

Conclusion

The assessment of Internet usage can be helpful in the treatment of psychological problems among adolescents. In a similar vein, for the treatment of Internet addicted adolescents, the evaluation of psychological distress, particularly depression and anxiety, must be considered. Effective counseling and intervention programs need to be designed and implemented for teaching healthy, moderate and non-addictive internet use for improving the overall psychological well-being of the adolescents.