

CASE REPORT

Extragastrointestinal Stromal Tumor Arising in the Pancreas: A Case Report with a Review of the Literature

Somanath Padhi¹, Ravikanth Kongara², Shantveer G Uppin¹, Megha S Uppin¹,
Aruna K Prayaga¹, Sundaram Challa¹, Bheerappa Nagari², Shastry A Regulagadda²

Departments of ¹Pathology and ²Surgical Gastroenterology, Nizam's Institute of Medical Sciences.
Panjagutta, Hyderabad, Andhra Pradesh, India

ABSTRACT

Context Extragastrointestinal stromal tumors arising in the pancreas are extremely rare. To date, only eight cases have been reported in the literature. **Case report** A 42-year-old female patient presented with gradually increasing abdominal pain of 6-month duration. Computerized tomography scan of the abdomen demonstrated a solid cystic mass in the body and tail of the pancreas. En-block R0 resection of the mass with distal pancreatectomy, splenectomy and left hemicolectomy was carried out following a radiological diagnosis of a malignant cystic neoplasm of the pancreas. Histopathological and immunohistochemical findings of the lesion were consistent with a gastrointestinal stromal tumor. **Conclusion** Extragastrointestinal stromal tumor of the pancreas, though rare, should be considered in the differential diagnosis of the more common cystic lesions at this site.

INTRODUCTION

Extragastrointestinal stromal tumors, by definition, originate from the soft tissues of the abdomen and retroperitoneum but display no connection to the wall or the serosal surface of the viscera [1]. These are known to arise from various anatomic sites such as the omentum, mesentery, retroperitoneum and gall bladder [1, 2]. Extragastrointestinal stromal tumors arising in the pancreas are extremely rare and only eight cases have been reported in the literature [3, 4, 5, 6, 7, 8, 9, 10]. We report a case of a pancreatic extragastrointestinal stromal tumor in a 42-year-old female patient together with a review of the literature.

CASE REPORT

A 42-year-old female patient presented with vague abdominal pain which had gradually increased in severity over a six month period. It was associated with loss of appetite and weight. The abdominal examination revealed an ill defined, large lobulated, firm mass of approximately 30x25 cm in the

epigastrium, left hypochondrium and extending up to left iliac fossa. The mass was non-ballotable and did not move with respiration.

Routine laboratory investigations were within normal limits except for anemia (hemoglobin: 8 g/dL; reference range: 12-15 g/dL). A contrast-enhanced computerized tomogram of the abdomen (Figure 1) revealed a large heterogeneously enhancing mass of mixed echogenicity (mostly cystic with a few solid areas), measuring 35x30x25 cm, in the region of the body and the tail of the pancreas. The mass did not communicate with the main pancreatic duct. There were no ascites or lymphadenopathy. On magnetic resonance imaging, the mass was hypointense on T1-weighted imaging and hyperintense (cystic) on T2-weighted imaging with hemorrhagic and calcific foci




Figure 1. Contrast-enhanced computerized tomogram of the abdomen: axial (a.) and coronal (b.) sections showing a large lobulated heterogeneously enhancing mixed echogenic lesion in the region of the body and the tail of the pancreas.

Received January 13th, 2010 - Accepted March 1st, 2010

Key words Gastrointestinal Stromal Tumors; Pancreas; Proto-Oncogene Proteins c-kit

Correspondence Sundaram Challa

Department of Pathology, Nizam's Institute of Medical Sciences,
Panjagutta, Hyderabad, Andhra Pradesh-500082, India
Phone: +91-40.2348.9051; Mobile: +91-984.949.6333
Fax: +91-40.2331.0076

E-mail: challa_sundaram@yahoo.com

Document URL <http://www.joplink.net/prev/201005/15.html>

(Figure 2). The possibility of a malignant cystic neoplasm of the pancreas was considered. Upper gastrointestinal endoscopy and colonoscopy showed no abnormalities. At laparotomy, a large mass was found to arise from the body and tail region of the pancreas. There was no attachment to the stomach, and the colon was stretched over the mass. There was no evidence of peritoneal dissemination. To achieve a R0 resection, a distal pancreatectomy, splenectomy and left hemicolectomy were performed.

A large lobulated solid and cystic mass measuring 35x30x25 cm with compressed pancreatic tissue at the periphery was seen at gross examination (Figure 3a). The mass was easily separated from colon. The cut section revealed areas of cystic degeneration with a blood clot (Figure 3b). Cystic fluid analysis for carcinoembryonic antigen (CEA) was within the normal range.

Histopathological examination of the specimen revealed a cellular lesion with compressed pancreatic tissue at the periphery. The cells showed both spindle and epithelioid morphology alternating with acellular hyalinized and myxoid stroma with thick-walled blood vessels. Cellular areas showed spindle cells arranged in fascicles. They had elongated nuclei with tapered ends




Figure 2. Magnetic resonance imaging showing the mass to be hypointense on T1-weighted imaging (a.) and hyperintense on T2-weighted imaging (b.).




Figure 3. a. Gross specimen showing a large lobulated mass in relation to the body and tail of the pancreas. **b.** Cut section of the mass showing cystic degeneration.

and moderate amount of cytoplasm (Figure 4). Mitotic figures were 6-8/50 high power field (HPF). Immunohistochemistry with CD117 (prediluted;




Figure 4. a. Section showing a spindle cell lesion with the adjacent pancreas. **b.** Fascicles of spindle cells separated by myxoid stroma.




Figure 5. Immunohistochemistry findings. **a.** CD117 positive in tumor cells. **b.** CD34 positive in tumor cells. **c.** Smooth muscle actin negative in tumor cells. **d.** S-100 negative in tumor cells.

BioGenex, Cherlapally, Hyderabad, India) showed intense membrane and cytoplasmic positivity in the cells (Figure 5a). The lesional cells were also positive

for CD34 (prediluted, BioGenex, Cherlapally, Hyderabad, India) (Figure 5b) but negative for smooth muscle actin (SMA) (dilution: 1:75; BioGenex, Cherlapally, Hyderabad, India) and S-100 (Figure 5cd). There was no infiltration into the omentum, colon or spleen. The large size (greater than 10 cm), location and mitotic activity of the 6-8/50 HPF classified the lesion as high risk [11]. The patient was kept on close follow-up without any postoperative chemotherapy.

DISCUSSION

The concept of a gastrointestinal stromal tumor (GIST) has recently been established, due to progress in methods of immunohistochemical analysis. These tumors express CD117 (c-KIT protein) which is expressed in the interstitial cells of Cajal and, thus, in animal investigations, the cells of origin of stromal tumors were recently suggested to be the interstitial cells of Cajal [12, 13]. In 2004, Yamamoto *et al.* [14] reported that extragastric stromal tumors share similar *KIT* mutations to typical GISTs which suggest that these tumors have a similar origin. The origin of

Table 1. Comparison of clinicopathological findings, treatment and follow-up data of reported cases of pancreatic gastrointestinal stromal tumors.

Case	Authors, Year	Age, Sex	Presentation	Location	Pathology	Treatment	Follow-up
#1	Neto <i>et al.</i> , 2004 [3]	67 years Female	Epigastric pain, bloating, weight loss	Body and tail	Size: 20 cm Solid cystic mass with necrotic foci Mitotic count: 120/50 HPF IHC: CD117 (+), CD34 (+); CK7 (-), CK20 (-), desmin (-), synaptophysin (-)	Distal pancreatectomy and splenectomy Imatinib mesylate	Recurrence and peritoneal dissemination 1 month after surgery
#2	Yamaura <i>et al.</i> , 2004 [4]	54 years Female	Incidental finding	Tail	Size: 14 cm Solid mass with cystic degeneration Few mitoses IHC: vimentin (+), CD34 (+); CD117 (-), SMA (-), S-100 (-)	Distal pancreatectomy, splenectomy, and partial gastrectomy	NED at 30 months
#3	Krska <i>et al.</i> , 2005 [5]	38 years Female	Abdominal pain	Body and head	Size: 17 cm Mitotic count: 1/50 HPF IHC: vimentin (+), CD34 (+); actin (-), CD117 (-), S-100 (-)	Partial pancreatectomy	NED at 30 months
#4	Daum <i>et al.</i> , 2005 [6]	70 years Female	Incidental finding	Head	Size: 10 cm Mitotic count: 2/50 HPF IHC: vimentin (-), CD34 (-), actin (-), CD117 (-), desmin (-), S-100 (-)	Whipple procedure, imatinib	NED at 6 months
#5	Showalter <i>et al.</i> , 2008 [7]	72 years Female	Incidental finding	Tail	Size: 7 cm Solid mass with necrotic foci Mitotic count: 3/50 HPF IHC: CD117 (+) SMA (-), S-100 (-)	Laparoscopic distal, pancreatectomy, and splenectomy	NED at 27 months
#6	Yan <i>et al.</i> , 2008 [8]	47 years Male	Nausea and vomiting	Uncinate process	Size: 2.4x2.1 cm Solid mass with necrotic foci Mitotic count: 3/50 HPF IHC: CD117 (+) SMA (-), S-100 (-)	N/A	N/A
#7	Haridhanavudi <i>et al.</i> , 2008 [9]	63 years Female	Fatigue and weakness	Body	Size: 16x11 cm Hemorrhagic cyst Mitotic count: less than 5/50 HPF IHC: CD117 (+), CD34 (+), SMA (+), MSA (+)	Cyst drainage and cystojejunostomy. Patient refused surgery and imatinib therapy	N/A
#8	Trabelsi <i>et al.</i> , 2009 [10]	52 years Female	Epigastric pain	Head	Size: 10.5x8x3 cm Mitotic count: 6/50 HPF IHC: CD117 (+), CD34 (+) SMA (-), S-100 (-), CK (-), synaptophysin (-)	Hemipancreaticoduodenectomy with antrectomy and partial colectomy	NED at 10 months
9	Present case	42 years Female	Abdominal pain, loss of weight and appetite	Body and tail	Size: 35x30x25 cm Mitotic count: 6-8/50 HPF IHC: CD117 (+), CD34 (+) SMA (-), S-100 (-), CK (-)	Distal pancreatectomy, splenectomy, and left hemicolectomy	NED at 10 months

(+): positive; (-): negative; CK: cytokeratin; IHC: immunohistochemistry; HPF: high power field; MSA: muscle specific actin; N/A: not available; NED: no evidence of disease; SMA: smooth muscle actin

stromal tumors in the pancreas with a similar expression of c-KIT protein is extremely rare and has been a matter of recent research.

The clinicopathological features and treatment outcomes of previously described pancreatic GISTs, including the present case, are presented in Table 1. There was a distinct female predominance with age ranging from 38 to 72 years (mean age 55 years). The majority occurred in the body and tail of the pancreas with an average size of 12 cm. Five of the eight cases had symptoms of abdominal pain with or without fatigue and weight loss. The mitotic rate in most of these tumors was low except in case 1 (120/50 HPF). This patient had rapid recurrence of the disease with peritoneal and nodal metastasis following surgery and imatinib therapy. For the first time, Yan *et al.* [8] and Harindhanavudhi T *et al.* [9] diagnosed two cases of pancreatic GISTs by means of endoscopic ultrasound guided fine needle aspiration cytology (EUS-FNA). On immunohistochemistry, five of the eight cases (cases #1, #5, #6, #7 and #8) demonstrated intense immunoreactivity for CD117 and CD34 whereas three cases (case #2, #3 and #4) showed negative staining for CD117. Reactivity for S-100 and SMA were variable in the lesional cells. Our patient had a relatively large tumor in comparison to other cases. Characteristic CD117 positivity was consistent with a diagnosis of GIST. The follow-up period was uneventful following definitive surgery without any evidence of metastasis.

The existence of interstitial cells of Cajal in the pancreas is difficult to explain. Popescu *et al.* [15] have recently demonstrated the existence of interstitial cells of Cajal in the human exocrine pancreas which have a phenotype similar to the enteric interstitial cells of Cajal. Moreover, Yamaura *et al.* [4] have shown the presence of c-KIT positive interstitial cells surrounding the intercalated ducts and acinus in the pancreas. Although the exact function of these cells is not clear, the discovery of pancreatic interstitial cells of Cajal supports the diagnosis of an extragastrointestinal stromal tumor arising solely from the pancreas.

The cytologic differential diagnoses of spindle cell proliferation in the pancreas include leiomyoma, leiomyosarcoma, GIST, fibromatosis, inflammatory fibroid polyp, schwannoma and gastrointestinal muscle sampling [8]. Immunohistochemical positivity of CD117 confirms the diagnosis of GIST. Cases of fibromatosis reported in the literature have been shown to react with an antibody directed against CD117, although this does not typically involve the pancreas [16].

The follow-up data of the previous case reports of pancreatic GISTs, including the present one, suggest that pancreatic stromal tumors may follow a benign course following definitive surgery as compared to extragastrointestinal stromal tumors arising from other sites.

To conclude, we presented a rare case of pancreatic GIST. The patient has been followed up postoperatively for ten months and there is no evidence

of recurrence or dissemination to date. Although rare, GISTs should be considered in the differential diagnosis of the more common cystic neoplasms of the pancreas. To clarify the relationship between pancreatic stromal tumors and the expression of c-KIT protein, additional research with a large number of cases of pancreatic stromal tumors having c-KIT protein expression is needed.

Conflict of interest The authors have no potential conflict of interest

References

- Weiss SW, Goldblum JR. Extra gastrointestinal stromal tumors. In: Enzinger and Weiss's Soft Tissue Tumors; fifth edition. Mosby Elsevier, 2008; p565-79. [ISBN 978-0-323-04628-2]
- Ortiz-Hidalgo C, de Leon Bojorge B, Albores-Saavedra J. Stromal tumor of the gallbladder with phenotype of interstitial cells of Cajal: a previously unrecognized neoplasm. *Am J Surg Pathol* 2000; 24:1420-3. [PMID 11023105]
- Neto MR, Machuca TN, Pinho RV, Yuasa LD, Bleggi-Torres LF. Gastrointestinal stromal tumor: report of two unusual cases. *Virchows Arch* 2004; 444:594-6. [PMID 15118853]
- Yamaura K, Kato K, Miyazawa M, Haba Y, Muramatsu A, Miyata K, Koide N. Stromal tumor of the pancreas with expression of c-kit protein: report of a case. *J Gastroenterol Hepatol* 2004; 19:467-70. [PMID 15012791]
- Krška Z, Pesková M, Povýsil C, Horejs J, Sedláčková E, Kudrnová Z. GIST of pancreas. *Prague Med Rep* 2005; 106:201-8. [PMID 16315768]
- Daum O, Klecka J, Ferda J, Treska V, Vanecek T, Sima R, et al. Gastrointestinal stromal tumor of the pancreas: Case report with documentation of KIT gene mutation. *Virchows Arch* 2005; 446:470-2. [PMID 15756592]
- Showalter SL, Lloyd JM, Glassman DT, Berger AC. Extragastric stromal tumor of the pancreas: case report and a review of the literature. *Arch Surg* 2008; 143:305-8. [PMID 18347279]
- Yan B M, Pai R K, Dam J V. Diagnosis of pancreatic gastrointestinal stromal tumor by EUS guided FNA. *JOP. J Pancreas (Online)* 2008; 9:192-6. [PMID 18326928]
- Harindhanavudhi T, Tanawuttiwat T, Pyle J, Silva R. Extra-Gastrointestinal Stromal Tumor Presenting as Hemorrhagic Pancreatic Cyst Diagnosed by EUS-FNA. *JOP. J Pancreas (Online)* 2009; 10:189-91. [PMID 19287116]
- Trabelsi A, Yacoub-Abid L B, Mtimet A, Ben Abdelkrim S, Hammedi F, Ben Ali A, Mokni M. Gastrointestinal stromal tumor of the pancreas: A case report and review of the literature. *North Am J Med Sci* 2009; 1:324-6.
- Reith JD, Goldblum JR, Lyles RH, Weiss SW. Extragastric (soft tissue) stromal tumors: an analysis of 48 cases with emphasis on histologic predictors of outcome. *Mod Pathol* 2000; 13:577-85. [PMID 10824931]
- Lecoin L, Gabella G, Le Douarin N. Origin of the c-kit positive interstitial cells in the avian bowel. *Development* 1996; 122:725-33. [PMID 8631250]
- Burns AJ, Herbert TM, Ward SM, Sanders KM. Interstitial cells of Cajal in the guinea-pig gastrointestinal tract as revealed by c-kit immunohistochemistry. *Cell Tissue Res* 1997; 290:11-20. [PMID 9377631]
- Yamamoto H, Oda Y, Kawaguchi K, Nakamura N, Takahira T, Tamiya S, et al. C-kit and PDGFRA mutations in extragastric stromal tumor (gastrointestinal stromal tumor of the soft tissue). *Am J Surg Pathol* 2004; 28:479-88. [PMID 15087667]

15. Popescu LM, Hinescu ME, Ionescu N, Ciontea SM, Cretoiu D, Ardelean C. Interstitial cells of Cajal in pancreas. *J Cell Mol Med* 2005; 9:169-90. [PMID 15784175]

16. Miettinen M. Are desmoid tumors kits positive? *Am J Surg Pathol* 2001; 25:549-50. [PMID 11257638]
