



## Habitual Routines and Their Role in Triggering Relapse Behavior

Olivia Grant\*

*Department of Behavioral Medicine, Northgate University, Edinburgh, United Kingdom*

### DESCRIPTION

Habitual Routines play a significant role in shaping behavior and in the context of addiction, they often become closely linked to substance use. Over time, repeated actions form patterns that are performed with little conscious thought. These patterns can act as powerful triggers for relapse, particularly when they are deeply embedded in daily life. Understanding how routines influence behavior is essential for developing effective strategies to maintain recovery (1). Habits are formed through repetition and reinforcement. When a behavior is performed repeatedly in a consistent context, it becomes associated with specific cues. For example, an individual who uses a substance after work each day may develop a routine in which the end of the workday automatically triggers the urge to use. This association becomes ingrained, making it difficult to separate the cue from the behavior. During recovery, these habitual patterns do not disappear immediately (2). Even after substance use has stopped, the cues associated with previous routines can continue to trigger cravings. This is because the brain has learned to expect a certain response in a given context. Breaking these patterns requires conscious effort and the introduction of alternative behaviors. One effective approach to addressing habitual triggers is to modify daily routines. This may involve changing the sequence of activities, altering the environment or introducing new behaviors at times when substance use previously occurred. For instance, replacing a routine of visiting a bar after work with attending a fitness class or engaging in a hobby can help disrupt the association between the cue and the behavior (3).

Consistency is important when establishing new routines. Repeated engagement in alternative activities strengthens new associations, gradually reducing the influence of old patterns. Over time, these new behaviors can become

automatic, replacing the previous habit. This process requires patience, as habit formation is gradual and influenced by repeated practice. Awareness is a critical component in managing habitual triggers. Individuals must recognize the cues that initiate their routines and understand how these cues influence behavior. Keeping a record of daily activities and noting moments of craving can help identify patterns that may not be immediately obvious (4,5). This information can then be used to develop targeted strategies for change. Environmental design also plays a role in shaping routines. Small changes in the physical environment can influence behavior by altering cues. For example, removing objects associated with substance use or rearranging living spaces can reduce exposure to triggers. Creating an environment that supports positive behaviors makes it easier to maintain new routines (6). Time-based triggers are another aspect of habitual behavior. Certain times of day may be associated with substance use due to past routines. These temporal cues can evoke cravings even in the absence of other triggers. Planning structured activities during these times can help redirect attention and reduce the likelihood of relapse.

Stress and fatigue can weaken the ability to resist habitual behaviors. When individuals are tired or under pressure, they may revert to automatic patterns. This highlights the importance of maintaining overall well-being, including adequate rest and stress management (7). By supporting physical and mental health, individuals can strengthen their ability to maintain new routines. Support from others can enhance efforts to change habitual behavior. Sharing goals with friends, family or support groups creates accountability and provides encouragement. Others can also offer suggestions for alternative activities and help reinforce positive changes. Technology can assist in managing routines by providing reminders and tracking progress (8,9). Applications that monitor daily activities can help individuals

**Received:** 24-Nov-2025; Manuscript No: IPJABT-25-23837; **Editor assigned:** 27-Nov-2025; PreQC No: IPJABT-25-23837 (PQ); **Reviewed:** 11-Dec-2025; QC No: IPJABT-25-23837; **Revised:** 18-Dec-2025; Manuscript No: IPJABT-25-23837 (R); **Published:** 24-Dec-2025 DOI: 10.35841/ipjabt-9.4.75

**Corresponding author:** Olivia Grant, Department of Behavioral Medicine, Northgate University, Edinburgh, United Kingdom; Email: olivia.grant@northgatemed.uk

**Citation:** Grant O (2025) Habitual Routines and Their Role in Triggering Relapse Behavior. J Addict Behav Ther. 9:75.

**Copyright:** ©2025 Grant O. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited

stay consistent with new behaviors. Feedback from these tools can reinforce progress and highlight areas that may require adjustment.

## CONCLUSION

In conclusion, changing habitual routines is a central aspect of recovery from addiction by understanding how habits are formed and how they influence behavior, individuals can take active steps to replace old patterns with healthier alternatives (10). This process involves awareness, consistency and support, all of which contribute to reducing the impact of behavioral triggers and sustaining long-term change.

## REFERENCES

1. Hodgins DC, El-Guebaly N (2004). Retrospective and prospective reports of precipitants to relapse in pathological gambling. *JCCP*.72(1):72.
2. Langham E, Thorne H, Browne M, Donaldson P, Rose J, et.al. (2015). Understanding gambling related harm: A proposed definition, conceptual framework, and taxonomy of harms. *BMC public health*. 16(1):80.
3. Ledgerwood DM, Petry NM (2006). What do we know about relapse in pathological gambling?. *Clin Psychol Rev*. 26(2):216-228.
4. Hodgins DC, el-Guebaly N (2010). The influence of substance dependence and mood disorders on outcome from pathological gambling: Five-year follow-up. *J Gambl Stud*. 26(1):117-127.
5. Daughters SB, Lejuez CW, Strong DR, Brown RA, Breen RB,et.al (2005). The relationship among negative affect, distress tolerance, and length of gambling abstinence attempt. *J. Gambl. Stud*. 21(4):363-378.
6. Moon SJ, Lee H (2020). Relapse to substance use: A concept analysis. *Nurs Forum*. 55(3):523-530.
7. Barnett BS, Parker SE, Weleff J (2022). United States National Institutes of Health grant funding for psychedelic-assisted therapy clinical trials from 2006–2020. *Int J Drug Policy*. 99:103473.
8. Fishman M, Wenzel K, Scodes J, Pavlicova M, Lee JD,et.al. (2020). Young adults have worse outcomes than older adults: Secondary analysis of a medication trial for opioid use disorder. *J Adolesc Health*. 67(6):778-785.
9. Sewak R, Spielholz NI (2018). Relapse prevention: Using sound to reduce the probability of recidivism and suffering following detoxification. *Med Hypotheses*. 118:84-91.
10. Rong C, Jiang HF, Zhang RW, Zhang LJ, Zhang JC, et.al. (2016). Factors associated with relapse among heroin addicts: evidence from a two-year community-based follow-up study in China. *IJERPH*. 13(2):177.