



Mind, Behaviour and Care: A Contemporary View of Psychiatry

Helena Strauss*

Department of Public Health Nutrition, Westbridge University, Berlin, Germany

DESCRIPTION

Psychiatry is a medical field concerned with the understanding, diagnosis and management of mental conditions that affect thinking, feeling and behaviour. It draws from medicine, psychology, neuroscience and social sciences to form a broad view of human mental function. The practice involves recognizing patterns of distress, identifying contributing factors and applying treatments that aim to improve well-being and daily functioning. Unlike many other areas of medicine, psychiatry often deals with experiences that are not directly visible, such as thoughts, emotions and perceptions, making careful communication and observation essential. Mental conditions can vary widely in form and severity. Some individuals may experience persistent sadness or lack of interest, commonly associated with depressive disorders, while others may face intense worry or fear linked to anxiety-related conditions. There are also illnesses that involve altered perception of reality, such as schizophrenia, where hallucinations or false beliefs may occur. Each condition presents a unique combination of symptoms and no two individuals experience them in exactly the same way. This variation requires clinicians to approach each case with flexibility and attention to personal context.

Assessment in psychiatry often begins with a detailed conversation. A clinician gathers information about current concerns, past experiences, medical history and social background. This process, sometimes called a psychiatric interview, helps in forming an initial impression(1,2). Observations of speech, mood, thought patterns and behaviour are combined with standardized tools to support diagnosis. While laboratory tests and brain imaging can assist in certain situations, most psychiatric diagnoses rely heavily on clinical evaluation (3,4). Biological factors play a significant role in many mental conditions. Brain chemistry, genetic

influences and physical health can all contribute to how a person thinks and feels. Neurotransmitters such as serotonin and dopamine are often discussed in relation to mood and perception. However, biological aspects alone do not fully explain mental illness. Psychological factors, including coping skills, personality traits and past experiences, are also important. Social influences such as relationships, cultural expectations and economic conditions further shape mental health outcomes (5). Psychiatry therefore adopts a biopsychosocial model, which considers all these dimensions together. Treatment approaches in psychiatry are diverse and often combined for better results. Medications are commonly used to manage symptoms. Antidepressants may help regulate mood, antipsychotics can reduce hallucinations or delusions and mood stabilizers assist in conditions involving emotional fluctuation. These medications require careful selection and monitoring to balance benefits and side effects. Psychotherapy is another major component of care (6). Techniques such as cognitive behavioral therapy help individuals identify and modify patterns of thinking and behavior that contribute to distress. Other forms of therapy focus on emotional expression, relationship patterns, or problem-solving skills.

In addition to medication and therapy, lifestyle factors are increasingly recognized as important in mental health care. Regular physical activity, adequate sleep, balanced nutrition and social interaction can all influence mental well-being. Clinicians often encourage patients to adopt healthy routines as part of a comprehensive treatment plan. Support from family and community also plays a vital role in recovery and stability. Stigma remains a challenge in the field of psychiatry (7). Misunderstandings about mental illness can lead to discrimination and reluctance to seek help. Many individuals delay treatment due to fear of judgment or lack of awareness. Public education and open discussion are important in

Received: 17-Mar-2025; Manuscript No: IPCP-26-23818; **Editor assigned:** 19-Mar-2025; PreQC No: IPCP -26-23818 (PQ); **Reviewed:** 02-Apr-2025; QC No: IPCP-26-23818; **Revised:** 09-Apr-2025; Manuscript No: IPCP-26-23818 (R); **Published:** 17-Apr-2025; DOI: 10.35248/2471-9854-11.02.59

Corresponding author: Helena Strauss, Department of Public Health Nutrition, Westbridge University, Berlin, Germany; Email: helena.strauss@westbridgeuni.de

Citation: Strauss H (2025). Mind, Behaviour and Care: A Contemporary View of Psychiatry. Clin Psychiatry. 11:59.

Copyright: © 2025 Strauss H. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

changing these attitudes. As awareness increases, more people are recognizing that mental health is an essential part of overall health and deserves the same attention as physical conditions (8).

Advances in neuroscience and technology continue to influence psychiatric practice. Brain imaging techniques provide insights into structure and function, while digital tools such as mobile health applications offer new ways to monitor symptoms and deliver interventions. Research into genetics and molecular biology is expanding understanding of risk factors and treatment response. Despite these developments, the human element remains central (9). Effective care depends on trust, empathy and clear communication between clinician and patient. Ethical considerations are also significant in psychiatry. Issues such as patient autonomy, informed consent and confidentiality require careful attention. In some cases, individuals may be unable to make decisions due to severe symptoms, raising questions about involuntary treatment. Clinicians must balance respect for individual rights with the need to ensure safety and appropriate care (10).

CONCLUSION

Advances in neuroscience and technology continue to influence psychiatric practice. Brain imaging techniques provide insights into structure and function, while digital tools such as mobile health applications offer new ways to monitor symptoms and deliver interventions. Research into genetics and molecular biology is expanding understanding of risk factors and treatment response. Despite these developments, the human element remains central. Effective care depends on trust, empathy and clear communication between clinician and patient.

REFERENCES

1. Dedoni S, Avdoshina V, Camoglio C, Siddi C, Fratta W, et al. (2022) K18- and CAG-hace2 transgenic mouse models and SARS-cov-2: Implications for neurodegeneration research. *Molecules*. 27(13):4142.
2. Vidal E, López-Figueroa C, Rodon J, Pérez M, Brustolin M, et al. (2022) Chronological brain lesions after SARS-cov-2 infection in hace2-transgenic mice. *Vet Pathol*. 59(4): 613-626.
3. Jeong GU, Lyu J, Kim KD, Chung YC, Yoon GY, et al. (2022) SARS-CoV-2 infection of microglia elicits proinflammatory activation and apoptotic cell death. *Microbiol Spectr*. 10(3):0109122.
4. Smeyne RJ, Eells JB, Chatterjee D, Byrne M, Akula SM, et al. (2022) COVID-19 infection enhances susceptibility to oxidative stress- induced parkinsonism. *Mov Disord*. 37(7):1394-1404.
5. Thompson JW, Blaine JD (1987) Use of ECT in the United States in 1975 and 1980. *Am J Psychiatry* 144(5): 557-562.
6. Eranti SV, McLoughlin DM (2003) Electroconvulsive therapy-state of the art. *Br J Psychiatry* 182: 8-9.
7. Hermann RC, Dorwart RA, Hoover CW, Brody J (1995) Variation in ECT use in the United States. *Am J Psychiatry* 152(6): 869-875. Ostergaard SD, Speed MS, Kellner CH, Mueller M, McClintock SM, et al. (2020) Electroconvulsive therapy (ECT) for moderate-severity major depression among the elderly: Data from the pride study. *J Affect Disord* 274: 1134-1141.
9. Kellner CH, Fink M, Knapp R, Petrides G, Husain M, et al. (2005) Relief of expressed suicidal intent by ECT: A consortium for research in ECT study. *Am J Psychiatry* 162(5): 977-982.
10. Bradvik L, Berglund M (2006) Long-term treatment and suicidal behavior in severe depression: ECT and antidepressant pharmacotherapy may have different effects on the occurrence and seriousness of suicide attempts. *Depress Anxiety* 23(1): 34-41.