

Review Article

Our Daily Use Chemical Products "The Enemy we Trust" Phthalates, Risks and Challenge to Avoid: An Update

Amr I M Hawal^{1*}, Malak A Alia², Fatma Elzahraa T Hussin³, Sherif M H Gad⁴, Gehan I M Mohamed⁵, Amr M M Amr⁶, Mohammad M Rizk Mohammad⁷

¹Department of Pediatrics and Neonatology, Amina Hospital, Ajman, UAE

²Department of Pediatrics, Prime Medical Center, Ajman, UAE

³Department of Pediatrics and Neonatology, Saudi-German Hospital, Ajman, UAE

⁴Department of Pediatrics and Neonatology, University Hospital Sharjah, Sharjah, UAE

⁵Depratment of Family Medicine, Primary Health Centre, Riyadh, Saudi Arabia

⁶Department of Medicine, Alexandria University, Alexandria, Egypt

⁷Department of Pediatrics and Neonatology, Mediclinic Al Ain Hospital, Al Ain, UAE

ABSTRACT

Background: The light on some chemical substances which are widely used in many and different aspects of our children daily practices and try to get the answers for the following important questions. Do these chemicals that our children are widely exposed to since even their prenatal period and throughout their lives have real high risks and dangers on their health? Is the use of these chemicals essential for our kids or we can avoid and replace them with other non-risky substances and hazardous chemical free products! Recently, a lot of systematic review studies have been done to investigate the possible relations between the most common chemical substances that we use daily and possible risks and health problems that may affect our children. Among the chemicals that have been studied intensively in the last few years are Phthalates which are known as endocrine disruptors due to their anti-androgenic and/or estrogenic effects!

The purpose and objective of our study: Are to provide an overview of some widely and commonly used chemicals that are introduced to many of the products essentially used by our children daily, and the possible relations between their use and some of the common health problems and disorders that affect children in order to raise the awareness among parents and health care providers regarding their potential health impacts on children and to provide a proper guidance that help to minimize the avoidable exposure to these risky chemicals and replace them with hazardous chemicals free products till we get enough studies that prove or disprove their risks and effects.

Peceived:	05 June 2022	Manuscript No:	IPPHR-23-16874
Receiveu.	05-June-2023	PreOC No:	IDDHR-23-16874 (DO)
Editor assigned:	09-June-2023	riede no.	IFFTIN-23-10874 (FQ)
Reviewed:	23-June-2023	QC No:	IPPHR-23-16874
Revised:	22-December-2023	Manuscript No:	IPPHR-23-16874 (R)
Published:	29-December-2023	DOI:	10.36648/2574-2817-8.4.35

Corrsponding author: Amr I M Hawal, Department of Pediatrics and Neonatology, Amina Hospital, Ajman, UAE; E-mail: amr106@gmail.com

Citation: Hawal AIM, Alia MA, Hussin FET, Gad SMH, Mohamed GIM, et al. (2023) Our Daily Use Chemical Products "The Enemy we Trust" Phthalates, Risks and Challenge to Avoid: An Update. Pediatr Heal Res. 8:35.

Copyright: © 2023 Hawal AIM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Discussion: Our study tries to search through, emphasis on and spot the light on the use of common chemical substances (Phthalates) that are widely used in most of our children life aspects, searches for their possible risks and hazards on our kids' health, tries to get relations and connections between these substances and common pediatric health problems and disorders. In our study we reviewed the data collected from many evidences based systematic reviews and cohort studies have dealt with these hazardous substances and stated the high incidence of their risks and their bad impacts on our kids' health and prove the relation of these chemicals to certain respiratory problems specially wheezing in childhood, ADHD, some atopic conditions and endocrine disruptions among children.

Conclusion: Our study concluded and suggested that it is better to eliminate exposure to these chemicals as evidences are rising against their safety. It's very important to increase the awareness among parents and health care providers regarding the possible risks of these chemicals and provide them the most proven practical tips for their avoidance. Further epidemiological studies should be conducted in the future to enhance our knowledge in this area.

Keywords: Pediatrics; Phthalates; Hazardous chemicals; Health

INTRODUCTION

Page 50

We live in a world of chemicals:

- Chemicals make our life easier, more convenient and more comfortable, but is it healthier?
- Since 1970's, more than 85000 chemicals have been approved for commercial use, yet only just one thousand of them have been examined and graded for their harmful potential.
- Most chemicals are used before being tested for safety and would only be regulated or banned after showing harmful effects on human, making us unwillingly experimental mices in this industrial world.

When we talk about chemicals, we don't only mean the pollution resulted from industry. We talk about things that we buy everyday presuming that if something is sold in supermarkets or pharmacies, and if everyone consume, then they should be safe. But the truth is bitter.

What is Phthalates?

- Phthalates are a group of man-made chemicals, added to Poly-Vinyl Chloride (PCV) to make this brittle plastic more flexible and durable (plasticizers).
- They were invented in 1920's and since then we witnessed a burst in plastic use globally, making us the third or fourth generation to be exposed to these chemicals [1].
- Many different phthalates exist with different properties, uses and health effects.
- High-molecular weight phthalates e.g. Di-2-Ethylhexyl Phthalate (DEHP) are primarily used as plasticizers in the manufacture of flexible vinyl plastic which, in turn, is used in: Waterproofs, raincoats and flip flop clothes.
- Domestic and construction: Flooring, wall coverings, shower curtains and hosepipes.
- Food contact applications like containers, food packaging and wraps.
- Toys, school supplies and childcare articles.
- Medical devices, like blood bags and tubing.

PVC products may contain up to 50% (by weight) phthalates and the more flexible the product is the more phthalate it contains [2].

 Low-molecular weight phthalates are used as solvents in personal-care products to hold fragrance and make products more effectively penetrate and moisturize the skin e.g. in perfumes, lotions, cosmetics, shampoos, aftershave, nail polishes, air fresheners and in lacquers, varnishes and coatings, including those used to provide timed releases in some pharmaceuticals.

LITERATURE REVIEW

How are we Exposed to Phthalates?

Since phthalates are not chemically bound to PCV polymers, they can leach, migrate or evaporate into indoor air, dust, food stuff and other materials. Human exposure to phthalates occurs through inhalation or ingestion of contaminated air and food as well as skin contact.

Phthalates in Processed and Packaged Food

- Processed and packaged food is the main source of human exposure.
- Mostly in fatty food, dairy products, processed cheese, processed meat, fish and oil, grain, ultra-processed food like soft drinks and fruit drinks, ready-to-eat-food and even in baby food and formulas [3].
- This contamination could happen either during processing or leaching from the packaging before consumer use.
- Phthalates exposures have also been associated with food venues, including fast food restaurants, school cafeteria lunches and dining out.

Daycare Products and Cosmetics

 Phthalates have proven to be very effective softeners in day care products like hair conditioners, hairsprays and body lotions so the use of certain personal care products especially makeup, lotion, emollients and sunscreens is associated with higher exposure to phthalates.

- Studies show that phthalates exist in the products when manufactured and their concentrations increase with the storage time indicating the possibility of leaching and migration of phthalates from the container also into the product.
- Phthalates made big news in 2008 when the American academy of pediatrics reported that infants exposed to infant-care products, specifically baby shampoos, baby lotions and baby powder, showed higher than normal levels of phthalates in their urine.
- This is also important in adolescent girls because they are in a critical age in hormonal development and have high use of personal care products.

Toys and Childcare Articles Made from PVC

- Plastic toys and childcare articles made from PVC are a great source of children exposure to phthalates as they can leach and contaminate the presumed safe surrounding environment or due to the hand-mouth habit in younger children [4].
- Soft toys contain higher levels of phthalates as compared to hard toys as primary function of phthalates is softening of hard plastic material.
- Because of their potential effects on human health, the European Union has banned the use of six specific phthalates in products for children.
- Toys and childcare articles made of PVC in other countries still contain variable levels of phthalates exceeding the EU regulations sometimes highly.
- If a toy is not designed to be sold on the European market by the manufacturer, it seems to be more likely to contain a phath phthalic acid ester.

Phthalates in Dust and Indoor Air

- Those dust particles accumulating on our floors might seem like a minor annoyance, but they could be exposing us to potentially harmful chemicals including phthalates.
- Studies confirm the occurrence of multiple phthalates sometimes in high concentrations in indoor dust, with DEHP being the most frequently detected phthalates in indoor dust [5].
- Humans are exposed to phthalates in dust *via* ingestion, inhalation and dermal uptake. Children are known to be more exposed to dust than adults due to their proximity to the floor and frequent hand-to-mouth behaviour.
- The age of the building, presence of foam mattresses and PVC flooring, presence of old soft plastic toys and the amount of plastic toys, the presence of electronic devices and the low cleaning frequency were associated with higher levels of phthalates in indoor dust [6].

latrogenic Phthalate Exposure

• Flexible plastic devices in hospitals like blood products bags, intravenous tubing, endotracheal tubes, feeding

tubes, catheter cannulas and others are most commonly made of Polyvinyl Chloride (PVC).

 In November of 2014, scientists at Johns Hopkins Bloomberg school of health reported that, "Because of phthalates in medical devices, premature babies admitted to the NICU can be exposed to 4,000 to 160,000 times the amount of phthalates considered safe.

Phthalates in Medication

- A lesser-known potential source of exposure is the timedrelease coatings on certain pharmaceuticals and dietary supplements.
- Phthalates are used also as excipients in numerous RX and OTC drug products and supplements from a wide range of therapeutic categories.
- Up to 50-fold higher levels of urinary phthalate metabolites have been observed in users of phthalate-containing drug products compared with non-users [7].
- Studies in children are lacking.

Phthalates Exposure

- Phthalates are rapidly metabolized and excreted in urine as a major elimination pathway, so the measurement of phthalate metabolites in urine is used to detect the recent exposure.
- Yet phthalate metabolites have been measured in various other body fluids, serum, semen, breast milk and saliva.
- Phthalates can cross the placental barrier and have been measured in amniotic fluid in human studies [8].
- The estimated daily exposure to one major phthalate, Di(2-Ethylhexyl) Phthalate (DEHP), ranges from 3 μg-30 μg/kg/d.
- According to the Forth national report on human exposure to environmental chemicals, nearly all Americans in every age group (age 6 or older) have detectable levels of phthalate metabolites and other endocrine disrupting chemicals in their urine [9].
- In Europe, according to the DEMOCOPHES project phthalate metabolites were detected in almost all participants (1844 mother-child pairs).
- In general, DEMOCOPHES results show that younger children (6-8 years) have higher exposure levels compared to older children (9-11 years), and higher levels in children compared to mothers, with the exception of MEP which is not regulated and is mainly used in cosmetics.
- A possible explanation is children's relatively higher intake: They are more exposed to dust, playing nearer the ground, have more frequent hand-to-mouth contact and they eat more than adults in relation to their weight.
- Consumption of convenience food, use of personal care products and indoor exposure to vinyl floors and wallpaper have all been linked to higher phthalate levels in urine.
- In Australia, data suggest that phthalate metabolites concentrations are at least two times higher than in the United States and Europe.

• A study in Saudi Arabia in 2017 measured major phthalate metabolites in urine samples of children aged 3-9 years [10-12].

The results showed that urinary levels of some phthalates were detected at higher levels than those reported previously in children worldwide, that was mostly related to the frequency of eating out, consumption of hamburgers, canned food, ice cream and the use of sun creams or sun spray.

Why Phthalates Raised Concern?

- Phthalates have received considerable attention because of their ubiquitous presence in the environment, frequent detection in human biomonitoring studies and demonstrated toxicity in rodents and humans.
- Phthalates are approved to have estrogenic effect making them Endocrine-Disrupting Chemicals (EDCs).

DISCUSSON

Page 52

Phthalates and Reproductive System

- Exposure to phthalates is of great concern, as they have been suspected to have anti-androgenic and estrogenic effects, which may alter the reproductive development of children.
- Animal studies indicate that gestational phthalate exposure is associated with adverse health outcomes, including disruption of endocrine and reproductive function and development.
- In females it induces pathological changes in ovaries, affects organ weights, disrupts estrous cyclicity, reduces fertility-related indices and causes some breeding complications.
- While in males, animal studies provide consistent evidence that certain phthalates target the developing reproductive system causing any genital anomaly. The connection is so clear the disorder is termed Phthalate Syndrome (PS) and the effects mirror a set of reproductive symptoms seen in human males, termed Testicular Dysgenesis Syndrome (TDS).
- In human, increased evidence shows that phthalates have adverse effects on Leydig cell development, thus causing lower androgen and Insulin-like 3 production and increasing the odds of any male newborn genital anomaly (termed Testicular Dysgenesis Syndrome (TDS) including cryptorchidism, hypospadias, shortened anogenital distance and isolated hydrocele.
- It also leads to disrupted gonadal function and spermatogenesis (low sperm counts and sperm motility and maturation) [13].
- And alters reproductive function of adolescent men including total testicular volume, reproductive hormones and semen volume.

Phthalates and Neurodevelopment

- We are witnessing an alarming increase in learning and behavioral problems in children.
- Although most of these problems are multifactorial, there are increased evidences that exposure to phthalate either prenatally or after birth can affect different aspects of neurodevelopment in children.
- It is strongly suggested to increase the odds of ADHD, attention deficit disorder, learning disability, social, thought, attention, cognitive and behavioral problems and may also affect the language development in children.
- The strongest evidence is for motor effects and BBP exposure in girls.

Phthalates have also been associated the development of:

- Overweight and obesity in school-age children.
- Increased BMI and waist circumference in adults and children.
- Increased insulin resistance, a precursor to diabetes, in teenager
- Childhood asthma.
- May worsen pulmonary function and airway inflammation in asthmatic children [14].

How to Avoid Phthalate Exposure

- We simply cannot escape our exposure to man-made chemicals, but there is a lot we can do to moderate and control our exposures.
- While onetime exposure to chemicals is often considered safe, many of these products are used every day and the effects of long-term exposure, especially when combined with other chemicals, is largely unstudied and unknown.

Awareness

Given the lax regulation of chemicals and the reactionary approach of governments' regulators, it is up to the consumer to be diligent about reading labels and making healthy choices to limit exposure to chemicals and toxins.

It is ideal to adopt the precautionary principle: "Until a chemical is found to be harmless, try to not use it", the precautionary principle means that you are maintaining awareness of what you are putting on and in your body and taking steps to avoid exposing yourself unnecessarily to toxins in your household and environment [15].

Minimize Plastic Use

- Limit exposure to plastics, especially anything with the number 3 or 7 on them.
- Use glass, ceramic, or metal containers for food and drink.
- If you have to use plastic, don't microwave it or wash it in the dishwasher to limit the leaching out of chemicals.
- Minimize plastic toys especially soft ones, instead you can use wooden toys or stuffed ones, Also, consider using silicone bottle nipple teethers and consider glass bottles for infants and babies that cannot feed themselves yet.

Avoid Food with High Concentration of Phthalates

- Avoid packaged foods: Studies have also shown that phthalates are particularly high in some packaged foods. Try to avoid these foods, especially for children.
- Foods that are higher in fat meats and cheeses, for instance are particularly prone to chemical leaching.
- Dairy items like heavy cream, ice cream and cheese had high to very high levels of phthalates.

CONCLUSION

Page 53

- Unfortunately, you will very rarely see phthalates listed on a product label. Luckily, there are clues. When it comes to cosmetics, the word "fragrance" or "perfume" on a label almost always means phthalates. What you want to see are claims like: "No synthetic fragrance" or "scented with only essential oils" or "phthalate-free." And always use only natural air freshener.
- Avoid personal care items where the scent is strong and lasts longer than 15-30 minutes as these products likely contain synthetic chemicals instead of pure essential oils.
- Minimize the use of unnecessary personal care products (The average man uses five to seven personal care products a day. The average woman uses nine to twelve products. The average teenage girl uses seventeen) and you can follow the rule (the less ingredients is better).
- Good resources for more information on the topic and safer alternatives include websites like EWG (ewg.org), breast cancer prevention partners (bcpp.o rg), IARC (iarc.fror monographs.iarc.fr), national institute of environmental health sciences (niehs.hih.gov), national cancer institute (cancer.gov) and the American cancer society (cancer.org).

REFERENCES

- 1. Wang Y, Zhu H, Kannan K (2019) A review of biomonitoring of phthalate exposures. Toxics. 5(2):7-21.
- Meeker JD, Sathyanarayana S, Swan SH (2009) Phthalates and other additives in plastics: Human exposure and associated health outcomes. Philos Trans R Soc B: Biol Sci. 364(1526):2097-2113.
- Broe A, Ennis ZN, PottegAard A, Hallas J, Ahern T, et al. (2017) Population exposure to phthalate-containing drugs. Basic Clin Pharmacol Toxicol. 121(3):153-158.
- Das MT, Ghosh P, Thakur IS (2014) Intake estimates of phthalate esters for South Delhi population based on exposure media assessment. Environ Pollut. 189:118-125.
- 5. Sakhi AK, Lillegaard IT, Voorspoels S, Carlsen MH, Loken EB, et al. (2014) Concentrations of phthalates and

bisphenol A in Norwegian foods and beverages and estimated dietary exposure in adults. Environ Int. 73:259-269.

- van Holderbeke M, Geerts L, Vanermen G, Servaes K, Sioen I (2014) Determination of contamination pathways of phthalates in food products sold on the Belgian market. Environ Res. 134:345-352.
- Beltifa A, Feriani A, Macherki M, Ghorbel A, Ghazouani L, et al. (2018) Persistent plasticizers and bisphenol in the cheese of Tunisian markets induced biochemical and histopathological alterations in male BALB/c mice. Environl Sci Pollut Res. 25:6545-6557.
- 8. Varshavsky JR, Morello-Frosch R, Woodruff TJ, Zota AR (2018) Dietary sources of cumulative phthalates exposure among the US general population in NHANES 2005-2014. Environ Int. 115:417-429.
- 9. Farooqi S, Rajendran K, Khanam R (2019) Analysis of phthalate esters in two different baby care products available in United Arab Emirates. Toxicol Mech Methods. 29(1):60-66.
- Berger KP, Kogut KR, Bradman A, She J, Gavin Q, et al. (2019) Personal care product use as a predictor of urinary concentrations of certain phthalates, parabens, and phenols in the HERMOSA study. J Expo Sci Environ Epidemiol. 29(1):21-32.
- Overgaard LE, Main KM, Frederiksen H, Stender S, Szecsi PB, et al. (2017) Children with atopic dermatitis and frequent emollient use have increased urinary levels of low-molecular-weight phthalate metabolites and parabens. Allergy. 72(11):1768-1777.
- 12. Ashworth MJ, Chappell A, Ashmore E, Fowles J (2018) Analysis and assessment of exposure to selected phthalates found in children's toys in christchurch, New Zealand. Int J Environl Res Public Health. 15(2):200.
- 13. Korfali SI, Sabra R, Jurdi M, Taleb RI (2013) Assessment of toxic metals and phthalates in children's toys and clays. Arch Environ Contam Toxicol. 65:368-381.
- 14. McCombie G, Biedermann S, Suter G, Biedermann M (2017) Survey on plasticizers currently found in PVC toys on the Swiss market: Banned phthalates are only a minor concern. J Environ Sci Health, Part A. 52(5):491-496.
- 15. Bi C, Liang Y, Xu Y (2015) Fate and transport of phthalates in indoor environments and the influence of temperature: A case study in a test house. Environ Sci Technol. 49(16):9674-9681.