

Proteomics Congress 2018: Sequential Double Bridging to Transplant with Diversified Anti-PD1 Monoclonal Antibodies retreatment in relapsed hodgkin lymphoma: A case report

Cinzia pellegrin

Santa Maria Delle Croci Hospital, Ravenna, Italy

Introduction :

Hodgkin lymphoma; Anti-PD1 monoclonal antibodies; Autologous stem cell transplant; Allogeneic transplantation; Retreatment Introduction Hodgkin lymphoma (HL) is the most common lymphoma affecting young population. Around 90% of early stage HL and 70% of advanced stage patients are cured with conventional chemotherapy, while 25% of patients experiences a relapse or results refractory to first line treatment. The latter subset of patients is a high priority for research and drug innovation. For patients with progression after first line therapy, the standard of care is a second line chemotherapy followed by autologous stem cell transplant (ASCT), with a 50% rate of cure.

Objective :

In this setting, the most favorable prognostic factor is achieving a complete remission (CR) before ASCT. For HL patients relapsing after ASCT, or with chemorefractory disease and/or ineligible for ASCT, there are several new agents such as brentuximab vedotin (BV), bendamustine, nivolumab and pembrolizumab that may improve their poor prognosis. In this setting allogeneic stem cell transplantation (alloSCT) is considered as a therapeutic option in young patients with a suitable donor and chemosensitive disease. Recent advances in the understanding of HL pathogenesis, interaction with tumor microenvironment and immune-escape mechanism have led to the identification of new therapeutic targets. Malignant Reed-Sternberg (RS) cells, which characterize HL, aberrantly express programmed death-1 (PD1) ligand and evade antitumor immune response by engaging programmed death-1 on immune effector cells. Pembrolizumab and nivolumab are programmed death-1 blockade (anti-PD1) monoclonal antibodies (mAb) that restore immunity against Hodgkin RS cells. Anti-PD1 have recently demonstrated high (60% to 80%) response rates in relapsed/refractory HL with manageable toxicities. For patients who responded to anti-PD1 and then relapsed, the choice ranges from aggressive regimens that enable alloSCT to palliative care for improving patient's quality of life. Here, we report on the first patient to our knowledge successfully re-treated with an anti-PD1 mAb prior to alloSCT.

Result :

Case Report In April 2014, a 21-year-old woman was diagnosed with classical HL involving lymphadenopathies both above and below the diaphragm, corresponding to a stage III B (pruritus). From April 2014 to October 2014 the patient received 6 cycles ABVD (adriamycin, bleomicine, dacarbazine, vinblastine). Final examination by 18F-fluorodeoxyglucose (FDG)-positron emission tomography (PET) and computed tomography (CT) demonstrated a residual metabolically active paramediastinal

lymphnodes. Subsequently, the patient received two cycles of IGEV (ifosfamide, gemcitabine, epirubicin and vinorelbine) regimen with successfully harvest of 20.4×10^6 CD34+ stem cells/kg after second cycle; the PET showed a progressive disease (PD), with increase of lymphadenopathies both above and below the diaphragm and appearance of multiple skeletal lesions. Due to symptomatic disease the patient was treated with six cycles of BV. Response assessment showed multifocal PD, which was histologically confirmed by laterocervical lymphadenectomy. The patient was treated with further therapy, namely bendamustine at 90 mg/m² on days 1 and 2 in a 28-day cycle schedule. She completed three courses of therapy, achieving a stable disease. In September 2015 the patient was enrolled in a phase II clinical trial and was treated with pembrolizumab at the dosage of 200 mg every 3 weeks with the aim to obtain a complete remission (CR) before being treated with ASCT. After 6 cycles of therapy the re-staging showed a CR and within one month the patient underwent ASCT.

Conclusion :

The chosen conditioning therapy was the BEAM (carmustine, etoposide, araC, melphalan) regimen. In September 2016, after six months from ASCT the patient had a recurrence of B symptoms (fever and pruritus). Disease relapse was documented by a routine follow-up PET which showed multiple hypermetabolic lymph nodes both above and below the diaphragm and multiple skeletal lesions. At this point, a search for a matched unrelated donor for alloSCT was started. Considering the disease chemorefractoriness but even the responsiveness to the anti-PD1 pembrolizumab, in October 2016 we decided to initiate an off-label therapy with another anti-PD1 mAb available in Italy, namely nivolumab at 3 mg/kg every 2 weeks, even in absence of retreatment clinical data. Treatment was well tolerated, B symptoms disappeared and patient clinical condition progressively improved over the following cycles. PET after 8 cycles showed a complete normalization of multiple hypermetabolic lymph nodes and skeletal lesions but also showed a new right inguinal lesion with SUV of 4.5. The CT scan (performed to check PET result) displayed small right inguinal lymph nodes hence it didn't confirm the presence of active disease. One month after the last course of nivolumab, in February 2017, following conditioning with cyclophosphamide (30 mg/kg), fludarabine (30 mg/m²), thiotepa (5 mg/kg) and antithymocyte globulin (6 mg/kg), the patient underwent transplant from an HLA-matched unrelated donor. The graft versus host disease (GVHD) prophylaxis was characterized by methotrexate (10 mg/m²) and cyclosporine 1 mg/kg from day -6 to -2 and 2 mg/kg from day -1. There was no development of acute or chronic post-transplant GVHD. The PET and CT re-staging

after 3, 6, 9 and 12 months from allo-SCT exhibited a CR status. Discussion In spite of generally favorable prognosis of HL, the prognosis of the primary refractory patients is poor. AlloSCT with reduced intensity conditioning, based on graft versus lymphoma, is a real chance to cure HL. The most favorable prognostic factor is the disease control before transplant, with the highest rate of cure associated to negative PET. PD1 signaling regulates immune response by decreasing T-cell activation and suppressing T-cell proliferation. In general, many clinical trials on anti-PD1 mAbs have demonstrated interesting results in lymphoma patients with chemorefractory disease. In particular, HL have a unique

biologic dependence on the PD1 pathway, because Hodgkin RS cells escape immune surveillance through a genetic alteration in chromosome 9p24.1, that leads to an overexpression of the PD1 ligands. Results from phase 2 pivotal trial of single agent nivolumab (3 mg/kg) administered every 2 weeks demonstrated an objective response rate (ORR) of 87% (17% CR) in relapsed/refractory HL patients and a manageable profile of toxicity. Recent reports have confirmed a durable response.

pierluigi.zinzani@unibo.it