



The Importance of Exposures in Culture-negative *Endocarditis* in an IV Drugs Abuser

Ensiyeh Rahimi^{1*}, Sara Gaderkhani¹, Arash Seifi¹, Mahsa Azadbakhsh Kanaforabi¹, Bahar Haghdoost², Amirhossein Eghbal³, Saharnaz Sazgarnejad⁴, Saber Esmaeili⁵

¹Department of Infectious Diseases and Topical Medicine, Tehran University of Medical Sciences, Iran

²National Institute of Health Research, Tehran University of Medical Sciences, Iran

³Students' Scientific Research Center, Tehran University of Medical Sciences, Iran

⁴School of medicine, Tehran University of Medical Sciences, Iran

⁵Department of Epidemiology and Biostatistics, Research Centre for Emerging and Reemerging infectious diseases, Iran

ABSTRACT

Background: Infective *Endocarditis* (IE) is a severe and life-threatening disease worldwide. Approximately 5% of IE cases have negative blood cultures. Risk factors are exposure to slow-growing bacteria and exposure to *Bartonella* species. Most patients have nonspecific symptoms such as fever, fatigue, and weight loss.

Case presentation: A 38 years old male with a history of addiction was referred to our center with severe dyspnea, high fever, chest pain, and shortness of breath. He went under CT angiography for pulmonary thromboembolism and TTE. According to the findings, the first diagnosis was right-side *endocarditis*, which led to septic embolism, so we started empiric antibiotic therapy with vancomycin plus ceftriaxone. After two weeks of constant fever despite broad-spectrum antibiotic therapy and pleural effusion drainage, we changed the antibiotic regimen to vancomycin plus meropenem, and re-sended blood cultures, which were negative 3 times, we considered Q fever, Bartonella, and Brucellosis as possible causes of culture-negative *endocarditis* according to the history of staying in the camp. We sent serology tests and blood PCRs to the laboratory of the Pasteur Institute of Iran. All of them were negative, but a serology test confirmed *Bartonella henselae* with a 1: 2048 titer. In the meantime, we changed the patient's drug regimen to doxycycline, plus rifampin, due to the high level of creatinine, and warfarin was switched to enoxaparin. Clinically, the patient recovered with no breathing difficulties or chest pain after two weeks of treatment, and the result of the *B. henselae* IgG follow-up serology was 1:512. Four weeks after admission, the patient was discharged from the hospital with instructions to see a cardiologist, quit abusing IV drugs, and continue taking doxycycline and enoxaparin.

Conclusion: The patient's social history regarding IV-drug abuse and low socioeconomic status are the most important considerations. Because of the increased risk factors in these patients, organisms such as *Bartonella Henselae*, *Coxiella Burnetii* and *Brucella* should also be investigated when cultures are negative.

Keywords: Culture-negative *endocarditis*; IV-drug abuse; low socioeconomic status; *Bartonella henselae*

INTRODUCTION

Infective *Endocarditis* (IE) is a severe and life-threatening disease worldwide [1]. However, because of non-specific

clinical presentations, confirmation and prescription of effective treatment are sometimes not simple tasks. History may help significantly in the diagnosis of IE; For example, it is more common among drug addicts, particularly Intravenous

Received:	04-June-2024	Manuscript No:	ipqpc-24-19859
Editor assigned:	06-June-2024	PreQC No:	ipqpc-24-19859 (PQ)
Reviewed:	20-June-2024	QC No:	ipqpc-24-19859
Revised:	25-June-2024	Manuscript No:	ipqpc-24-19859 (R)
Published:	02-July-2024	DOI:	10.36648/1479-1064.32.3.15

Corresponding author Ensiyeh Rahimi, Department of Infectious Diseases and Topical Medicine, Tehran University of Medical Sciences, Iran, E-mail: erij.yh@yahoo.com

Citation Rahimi E, Gaderkhani S, Seifi A, Kanaforabi MA, Haghdoost B, et al. (2024) The Importance of Exposures in Culture-negative *Endocarditis* in an IV Drugs Abuser. Qual Prim Care. 32:15.

Copyright © 2024 Rahimi E, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

drug users [2,3]. Approximately 5% of IE cases have negative blood cultures, and its risk factors are exposure to slow-growing bacteria such as *Bartonella* species, fastidious nonbacterial organisms, previous antibiotic use, underlying valvular heart disease, and intracardiac or vascular device or other foreign bodies in contact with the blood [4-6]. Most patients have nonspecific symptoms such as fever, fatigue, and weight loss. In a case series comprising 348 blood culture-negative *endocarditis* cases from France, almost all the patients had a fever as a presenting symptom. In contrast, about 50% to 70% had symptoms of heart failure, such as exertional dyspnea, and about 50% had insidious weight loss [7].

Bartonella species is a small, intracellular, gram-negative, and very fastidious rod mainly transmitted by vectors; they are the second most common cause of culture-negative *endocarditis*. Among the cases of *Bartonella endocarditis*, two species predominantly implicated in causing culture-negative *endocarditis* are *B. henselae* and *B. Quintana* [8]. These bacteria have been isolated from many mammalian species, including cats and dogs. It can cause mild infection to severe and life-threatening *endocarditis* in humans and dogs. In dogs, several *Bartonella* species have been identified; one of the most common ones is *B. henselae*. It should be mentioned that in recent years, more cases of culture-negative *endocarditis* have been reported from developing countries [9]. This report presents a case of a patient with culture-negative *B. henselae endocarditis* from Iran, diagnosed using a combined diagnostic approach that included clinical evaluation, imaging, epidemiology, serology, echocardiography, and Transthoracic Echocardiography (TTE).

CASE REPRESENTATION

Patient Information

The patient was a 38 years old male, single and unemployed with a history of addiction, who had received care from a treatment camp for 6 months. He was discharged when he was on methadone maintenance therapy just before his first hospital admission. He was an IV drug abuser with a history of regularly using amphetamine, cocaine, and heroin. He also smoked cigarettes for 20 years. He also exposed that he had unsafe sex and lost contact with dogs and cats.

Clinical Findings Timeline

The symptoms onset was 20 days before admission when he had fever and chills besides shortness of breath in the camp. The patient was hospitalized in another care center for five days as a suspected COVID-19 case and he was treated with Remdesivir, but his nasopharyngeal and oropharyngeal COVID-PCR tests were negative. Having been discharged from the hospital, he started using amphetamine again, which deteriorated his condition.

This time he was admitted to our center, as a referral center, with severe dyspnea, high fever (40°C), chills, chest pain, myalgia, and hemoptysis. The patient was ill and, on physical examination, he had tachycardia (Heart rate=107) and tachypnea (Respiratory rate=28), with low blood pressure (90/60), fluctuated oxygen saturation which was less than 92%, normal heart auscultation, no clubbing, no splenomegaly, and

no lymphadenopathy. **Table 1** summarizes the lab results.

Table 1: The results of main lab tests at the admission time

The results of main lab tests at the admission time	
WBC	18,100 (normal range between 4,500-11,000)
Procalcitonin	More than 30 (normal range between 0.5-5)
Covid PCR test	Negative
CRP	174 (normal range: Less than 10 mg/L)
Creatinine	1.1
Troponin	Negative
D-dimer	25.14 (normal range: Less than 0.50)
Blood culture (3 times)	Negative
HCV Ab	Positive
HCV PCR	Negative

A 65-year-old patient was referred for investigation of an axonal sensorimotor neuropathy that has been evolving for 5 months. Possible infectious and neoplastic causes were ruled out by examination, laboratory tests and imaging. The patient reported ocular and oral dryness. Minor salivary gland biopsy highlighted the presence of focal lymphocytic sialadenitis with a focus score of 1 foci/4 mm². Schirmer's test was inferior to 5 mm/5 min in both eyes. Laboratory investigations revealed anterior pituitary failure including secondary adrenal insufficiency, central hypothyroidism and hypogonadotropic hypogonadism. Magnetic resonance imaging of the hypothalamo-pituitary region demonstrated signs of autoimmune hypophysitis. During hospitalization, the patient developed anemia and thrombocytopenia with schizocytes count at 4%. The diagnosis of TMA associated to SS, without renal failure was made. The patient was commenced on intravenous immunoglobulin, corticosteroids, azathioprine and pregabalin. An improvement was noticed in her neuropathic pain, anemia and thrombocytopenia. Schizocytes count was reduced progressively.

Diagnostic Assessment and Therapeutic Intervention

Due to fever, dyspnea, and an SPO₂ below 90%, a Computed Tomography (CT) was performed. The chest CT showed moderate pleural effusion and bilateral emphysema in the apex of the lungs and cardiomegaly. Pelvic and abdominal ultrasounds were normal. Because of dyspnea, tachycardia elevated D-dimer, and SpO₂ of less than 90%, we decided to take CT angiography for Pulmonary Thromboembolism (PTE). CT angiography showed PTE, moderate pleural effusion in the right lung compatible with empyema, and a blurred wedge in the Left Upper Lobe (LUL). We put the patient under anticoagulation therapy with warfarin, after PTE was diagnosed, and based on the loculated pleural fluid in the CT scan, empyema pattern, and the patient's fever, the effusion was tapped, and a chest tube was placed. The patient's pleural fluid was analyzed, which was exudative with Alb=900, Pro=2400, LDH=1184, RBC=50, WBC=3400, Glucose=80, and negative culture. The lung field was found to have a septic embolism pattern, which led to an

echocardiogram. Upon echocardiography, small vegetation was detected; therefore, a Transthoracic Echocardiogram (TTE) was conducted to confirm. TTE showed severe tricuspid valve damage, 8 millimeter vegetation in place, increased pulmonary artery pressure (PAP=40), and Ejection Fraction (EF)=45%, confirming the vegetation on the atrial side of the tricuspid valve.

According to the findings, the first diagnosis was right-side *endocarditis*, which led to septic embolism, so we started empiric antibiotic therapy with vancomycin plus ceftriaxone. After 2 weeks of constant fever despite broad-spectrum antibiotic therapy and pleural effusion drainage, we changed the antibiotic regimen to vancomycin plus meropenem, and, re-evaluated our diagnosis by taking a detailed history, re-sending blood cultures, which were negative 3 times, performing abdominal ultrasound and investigating the possible causes of culture-negative *endocarditis* including Q fever, Bartonella, and brucellosis, according to the history of staying in the camp. The serology and blood PCRs were sent to the laboratory of the Pasteur Institute of Iran. The PCR and serology for Q-fever were negative. Serology wright and 2-ME test for brucellosis were negative, but a serology test confirmed for *Bartonella henselae* with a 1:2048 titer. In the meantime, we changed the patient's drug regimen to doxycycline (100 mg twice daily), instead of aminoglycosides, plus rifampin (300 mg twice daily), due to the high level of creatinine (3 mg/dL-3.3 mg/dL), and warfarin was switched to enoxaparin due to rifampin and warfarin drug interactions and patient noncompliance. After 72 hours he responded to this drug combination significantly; Fever, tachycardia, and dyspnea improved, creatinine levels decreased and the pleural fluid test result was negative for Bartonella. After two weeks, rifampin was discontinued, but he received an extended course of doxycycline monotherapy. Also, due to severe tricuspid valve insufficiency, he underwent medical treatment, but because of his addiction, the cardiac surgery service did not recommend him for valvoplasty. A summary of the clinical practice of the patient is shown in **Figure 1**.




Figure 1: Timeline of symptoms onset and interventions

Follow-up and Outcomes

After 2 weeks of treatment with doxycycline plus rifampin, clinically, the patient recovered with no difficulty in breathing and no chest pain. The results of the follow-up serology for *B. henselae* IgG were 1:512. The patient was discharged from the hospital approximately four weeks after admission, with a recommendation to visit a cardiologist, stop his IV drug abuse, and continue doxycycline and enoxaparin. A controlled chest X-ray showed improvement, and the chest tube was discontinued. The patient left the hospital in good

general condition, without fever, with a normal heart rate and acceptable SpO₂ (>94%).

RESULTS AND DISCUSSION

Blood culture-negative IE has a high variation in prevalence among countries; the reports vary from 2.5%-70% of all infective *endocarditis* based on zoonotic agent exposure, antibiotic prescription pattern, and diagnostic tools [10]. *Bartonella* species are among the main microorganisms in culture-negative *endocarditis*. Meanwhile, among all 45 known *Bartonella* species, *B. henselae* is the second most common cause of culture-negative *endocarditis* [11]. While cats and cat-scratch are considered the main reservoir of *B. henselae*, dogs could also transmit infection as accidental hosts [12]. Moreover, although *Bartonella endocarditis* is a worldwide issue and there are case reports from all parts of the world, most reported cases are from European countries and the Americas [11].

In this article, we are reporting a case of *Bartonella endocarditis* in Iran; it is notable that since *Bartonella endocarditis* is not common in Iran, usually physician does not put this agent on the list of their differential diagnosis, which causes several challenges to achieving a timely diagnosis and treatment of *Bartonella endocarditis*. Besides, the final diagnosis is even more challenging due to non-specific symptoms, which are the same as other subacute *endocarditis* or infective syndromes. Moreover, the diagnosis is more challenging during the COVID-19 pandemic and the similarity between the typical symptoms of SARS-CoV-2 and *endocarditis* [13]. Nevertheless, several diagnostic tools are available; despite their limitations, such as low specificity and cross-reactions, serology is still considered valuable in diagnosing *Bartonella endocarditis* [14]. Also, PCR testing on whole blood or plasma specimen, with approximate sensitivity of 58% and specificity of 100%, is an effective diagnostic tool [15].

In this case, the patient had a history of contact with dogs, but based on previous studies, connection with cats is more common in most *B. henselae endocarditis* cases. Moreover, in the current case, one of the first complaints of the patient was exertional dyspnea, and heart failure manifestations are considered an acute coronary syndrome. At the same time, based on the study conducted by Okaro, et al., approximately 50% to 70% of *Bartonella endocarditis* presents symptoms of heart failure, including exertional dyspnea [11]. Low socioeconomic status is one of the underlying determinants of developing *B. Quintana endocarditis*. While in our patient, who was diagnosed with *B. henselae*, the patient's social history regarding IV-drug abuse and low socioeconomic status were the most important considerations [11]. Also, prognosis and treatment must consider epidemiological exposure and alternate diagnoses for culture-negative *endocarditis* in individuals who don't respond to therapy.

CONCLUSION

The patient's social history regarding IV-drug abuse and low socioeconomic status are the most important considerations. Because of the increased risk factors in these patients, organisms such as *Bartonella henselae*, *Coxiella burnetii* and

Brucella should also be investigated when cultures are negative.

ETHICAL APPROVAL

This study has been approved by the ethics committee of Tehran University of Medical Sciences, adheres to the Declaration of Helsinki and informed consent was taken. The patient has given a written informed consent for the publication. The authors confirm that written consent has been obtained from the patient for submission and publication. All the patient's data and materials are provided in the manuscript.

ACKNOWLEDGEMENT

None.

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

REFERENCES

- Hogevik H, Olaison L, Andersson R, Lindberg J, Alestig K (1995) Epidemiologic aspects of infective *endocarditis* in an urban population. A 5-year prospective study. *Medicine*. 74(6):324-39.
- Yew HS, Murdoch DR (2012) Global trends in infective *endocarditis* epidemiology. *Curr Infect Dis Rep*. 14(4):367-72.
- Murdoch DR, Corey GR, Hoen B, Miró JM, Fowler VG, et al. (2009) Clinical presentation, etiology, and outcome of infective *endocarditis* in the 21st century: The International Collaboration on *Endocarditis*-prospective cohort study. *Arch Intern Med*. 169(5):463-73.
- Tunkel AR, Kaye D (1992) *Endocarditis* with negative blood cultures. *N Engl J Med*. 326(18):1215-7.
- Van SRE (1982) Culture-negative *endocarditis*. *Mayo Clin Proc*. 57(3):149-54.
- Katsouli A, Massad MG (2013) Current issues in diagnosing and managing blood culture-negative infective and non-infective *endocarditis*. *Ann Thorac Surg*. 95(4):1467-74.
- Houpikian P, Raoult D (2005) Blood culture-negative *endocarditis* in a reference center: Etiologic diagnosis of 348 cases. *Medicine*. 84(3):162-73.
- Patel R, Call M, Schnee A (2022) A case of *Bartonella henselae* native valve *endocarditis* presenting with crescentic glomerulonephritis. *IDCases*. 27:e01366.
- Chomel BB, Kasten R, Williams C, Wey A, Henn J, et al. (2009) *Bartonella endocarditis*: A pathology shared by animal reservoirs and patients. *Ann N Y Acad Sci*. 1166(1):120-6.
- Fournier PE, Gouriet F, Casalta JP, Lepidi H, Chaudet H, et al. (2017) Blood culture-negative *endocarditis*: Improving the diagnostic yield using new diagnostic tools. *Medicine (Baltimore)*. 96(47):e8392.
- Okaro U, Addisu A, Casanas B, Anderson B (2017) *Bartonella* species, an emerging cause of blood-culture-negative *endocarditis*. *Clin Microbiol Rev*. 30(3):709-46.
- Chomel BB, Boulouis HJ, Maruyama S, Breitschwerdt EB (2006) *Bartonella* spp. in pets and effect on human health. *Emerg Infect Dis*. 12(3):389-94.
- Arbune M, Iancu AV, Lupasteanu G, Vasile MC, Stefanescu V (2021) A Challenge of COVID-19: Associated infective *endocarditis* with *Streptococcus gordonii* in a young immunocompetent patient. *Medicina (Kaunas)*. 57(12):1298.
- Barka NE, Hadfield T, Patnaik M, Schwartzman WA, Peter JB (1993) EIA for detection of *Rochalimaea henselae* reactive IgG, IgM, and IgA antibodies in patients with suspected cat-scratch disease. *J Infect Dis*. 167(6):1503-4.
- Sanogo YO, Zeaiter Z, Caruso G, Merola F, Shpynov S, et al. (2003) *Bartonella henselae* in *Ixodes ricinus* ticks (Acari: Ixodida) removed from humans, Belluno province, Italy. *Emerg Infect Dis*. 9(3):329-32.