

Pelagia Research Library

Advances in Applied Science Research, 2013, 4(2):54-59



Thermo-acoustical molecular interaction studies in ternary liquid mixtures using ultrasonic technique at 303K

A. A. Mistry¹, V. D. Bhandakkar² and O. P. Chimankar³

¹Department of Physics, Anand Niketan College, Warora, Maharashtra ²Department of Electronics, Anand Niketan College, Warora, Maharashtra ³Department of Physics, R. T. M. Nagpur University, Nagpur

ABSTRACT

The ultrasonic studies in liquids are great use in understanding the nature and strength of molecular interaction. The thermo-acoustical parameters for ternary liquids mixtures namely, Chlorobenzene + Benzene + Tetrahydrofuran (THF) have been estimated, from the measured values of ultrasonic velocity (U), density (ρ) and viscosity (η). Using the measured data, some of the acoustical parameters such as, adiabatic compressibility (β_a) free length (L_f), free volume (V_f), internal pressure (π_i), relaxation time (τ), and Gibb's free energy (ΔG) are evaluated at the temperature 303K. The present paper represents the nonlinear variation of ultrasonic velocity and the thermo-acoustical parameters lead to dipole- induced dipole interaction between chlorobenzene and benzene molecule while dipole-dipole interaction between the chlorobenzene and THF molecule. The behavior of these parameters with composition of the mixture has been discussed in terms of molecular interaction between the components of the liquids.

Keywords: Ultrasonic velocity, acoustical parameters, molecular interactions Chlorobenzene, Benzene, Tetrahydrofuran (THF) and ternary mixtures.

INTRODUCTION

Ultrasonic study is very much useful for characterizing the Physico-chemical behavior of liquids mixtures and measurements are used to study molecular interactions in the liquids [1, 2]. The method of studying in molecular interaction from the knowledge of variation of acoustic parameters along with their excess values with change in mole fraction gives an insight into the molecular process [3, 4]. The increase or decreases in ultrasonic velocities have been employed in understanding the nature of molecular interaction in the pure liquids binary mixtures [5-8] and ternary mixtures [9, 10]. The study of liquid mixtures containing of polar and non-polar components find applications in industrial and technological process [11].

The mixing of different components gives rise to solutions that generally do not behave ideally [12, 13]. Further these properties have been widely used to study the molecular interaction between the various species in the mixture [14, 15].

In present study ultrasonic velocity, density and viscosity were measured experimentally for the ternary systems namely Chlorobenzene + Benzene + Tetrahydrofuran at 303K. From the measured data, thermo-acoustical parameters have been computed and the results are analyzed in the light of molecular interaction.

.....(2)

MATERIALS AND METHODS

The liquid mixtures of various concentrations in mole fraction were prepared by taking AR grade chemicals. In the ternary mixture system, the mole fraction of second component, Benzene (X_2) is kept fixed arbitrarily at $X_2 = 0.40$. Mole fraction of Chlorobenzene (X_1) is increased from 0.00 to 0.60 while mole fraction of THF (X_3) is decreased from 0.60 to 0.00.

The ultrasonic velocity in the liquid mixtures have been measured using an ultrasonic interferometer (Mittal type: Model: M-83) working at frequency 3 MHz with an overall accuracy of $\pm 0.1 \text{ ms}^{-1}$, an electronically digital operated constant temperature water bath has been used to circulate water through the double walled measuring cell made up of a steel containing the experimental solution at the desired temperature. The density of pure liquids and liquid mixtures was determined using a 25ml specific gravity bottle with an accuracy of $\pm 0.1 \text{ Kgm}^{-3}$. An Ostwald's viscometer was used for the viscosity measurement of pure liquids and liquid mixtures with accuracy 0.0001Nm⁻²s. The viscometer was calibrated before used. The time of flow of water (t_w) and time flow of solution (t_s) was measured with digital stop watch. All the precautions were taken to minimize the possible experimental error.

3. THEORY

• Adiabatic compressibility (β_a) has been calculated from the ultrasonic velocity (U) and the density (ρ) of the medium using the equation as:

$$\beta_{a} = 1 / U^{2} \rho$$
(1)

• Intermolecular free length has been determined as: $L_f = K_T (\beta_a)^{1/2}$

Where K_T – is a Jacobson's constant.

• Free volume in terms of ultrasonic velocity (U) and viscosity of the liquid (η) as :	
$V_{f} = [M_{eff.} U / K \eta]^{3/2}$	(3)

Where $M_{eff.}$ is the effective molecular weight ($M_{eff.} = \sum m_i X_i$, in which m_i and X_i are the molecular weights and the mole fraction of the individual constituents respectively). K is a temperature independent constant which is equal to 4.28 x 10⁹ for all liquids.

• Internal pressure is calculated by	
$\Pi_{i} = bRT (K \eta / U)^{1/2} (\rho^{2/3} / M^{7/6})$	(4)

Where, b stands for cubic packing which is assumed to be 2 for all liquids and K is a dimension less constant, independent of temperature and the nature of liquids and its value is 4.28×10^9 , T is the absolute temperature and M_{eff.} is the effective molecular weight.

• Relaxation time in terms of adiabatic compressibility (β_a) and viscosity of the liquid (η) as: $\tau = 4/3 \beta_a \eta$ (5)

• **Gibb's free energy** can be calculated from the following relation: $\Delta G = KT \log (KT\tau /h)$

Where, τ is the relaxation time, K the Boltzmann constant, T the absolute temperature and h is the Planks constant.

Table 1:- Comparison of experimental values of density (ρ) , viscosity (η) and ultrasonic velocity (U) of pure liquids at 303K with literature values:

Liquids	Density(ρ) (Kgm ⁻³)		Viscosity(η)(10 ⁻³ NSm ⁻²)		Velocity(U) (ms ⁻¹)	
	Expt.	Literature	Expt.	Literature	Expt.	Literature
Chlorobenzene	1025.00	1056.36	0.6995	0.7027	1220.50	1236.00
Benzene	852.10	865.70	0.6510	0.6040	1280.30	1285.00
Tetrahydrofuran	872.30	875.10	0.6389	0.6455	1235.20	1240.10

$Table 2:- The experimental values of density (\rho), viscosity (\eta) and velocity (U) for the ternary system- Chlorobenzene + Benzene + Tetrahydrofuran at 303K.$

Mole f	raction X ₂	Density (ρ)	Viscosity (η) (10 ⁻³ NSm ⁻²)	Velocity (U)
0.0000	0.6000	662.51	0.408	1284.00
0.1000	0.5000	854.67	0.534	1254.00
0.1999	0.4001	861.99	0.548	1233.00
0.2999	0.3001	896.77	0.580	1225.80
0.4000	0.2000	931.54	0.612	1215.60
0.5001	0.0999	933.37	0.619	1195.80
0.6000	0.0000	942.52	0.631	1180.80

 $Table \ 3:- \ The \ experimental \ values \ of \ adiabatic \ compressibility \ (\beta_a), free \ length \ (L_f) \ and \ free \ volume \ (V_f)) \ for \ the \ ternary \ system-Chlorobenzene + \ Benzene + \ Tetrahydrofuran \ at \ 303K.$

Mole f	raction	Compressibility (β_a)	Free length (L_f)	Free volume (V_f)
Λ_1	Λ_3	(10 m N)	(10°m)	(10 m mol)
0.0000	0.6000	9.155	0.6051	4.0529
0.1000	0.5000	7.440	0.5455	2.8237
0.1999	0.4001	7.630	0.5524	2.8607
0.2999	0.3001	7.421	0.5448	2.7930
0.4000	0.2000	7.264	0.5390	2.7209
0.5001	0.0999	7.492	0.5474	2.7926
0.6000	0.0000	7.609	0.5517	2.8378

Table 4:- The experimental values of internal pressure (π_i), relaxation time (τ) and Gibb's free energy (ΔG) for the ternary system-Chlorobenzene + Benzene + Tetrahydrofuran at 303K

Mole f	raction	Internal pressure (π_i)	Relaxation time (τ)	Gibb's free energy (ΔG) (10 ⁻²⁰ K Imple ⁻¹)
0.0000	0.6000	202.20	0.4082	
0.1000	0.5000	377.10	0.5304	0.5304
0.1999	0.4001	365.19	0.5575	0.5575
0.2999	0.3001	366.08	0.5746	0.5746
0.4000	0.2000	367.66	0.5935	0.5935
0.5001	0.0999	354.27	0.6188	0.6188
0.6000	0.0000	344.92	0.6402	0.6402

Graphs: - Fig. 1-9, shows variation of density (ρ), viscosity (η), ultrasonic velocity (U), adiabatic compressibility(β_a), free length (L_f), free volume(V_f), internal pressure(π_i), relaxation time(τ), and Gibb's free energy(ΔG) with respect to mole fraction at temperature 303K.



Fig.(2): Variation of Viscocity with mole fraction





RESULTS AND DISCUSSION

The experimentally measured and literature values of density, viscosity and Velocity for pure liquids at 303K are presented in **Table-1**. Experimental values of density, viscosity and Velocity for the ternary system namely, Chlorobenzene + Benzene + Tetrahydrofuran at 303K are given in **Table-2**. The thermodynamics parameters such as adiabatic compressibility (β_a), free length (L_f) free volume (V_f) internal pressure (π_i), relaxation time (τ) and Gibb's free energy (ΔG) at temperature 303K are listed in **Table-3&4**. The variation of density, viscosity and ultrasonic velocity Vs mole fraction for the ternary system- Chlorobenzene + Benzene + Tetrahydrofuran at 303K are shown in **fig.-1**, **2 and 3 respectively**. While other thermodynamics parameters such as adiabatic compressibility (β_a), free length (L_f) relaxation time (τ) and Gibb's free energy (ΔG) at temperature 303K are shown in **fig.-4,5,6,7,8 and 9 respectively**.

From the Table- 2 it is observed that, the density (ρ), viscosity (η) increases with increase in mole fraction for the system- Chlorobenzene + Benzene + Tetrahydrofuran and ultrasonic velocity decreases with increasing mole fraction for the system. The decreases in ultrasonic velocity are due to the decrease in adiabatic compressibility and free length of the liquid mixtures. This may lead to the presence of dispersive force (London force) between the molecules of the liquid mixture [16]. The adiabatic compressibility and intermolecular free length are the deciding factors of ultrasonic velocity in the ternary liquid mixtures. The decrease in free volume (fig.6) and increase in internal pressure (fig.7) represent that, the strength of the interaction decreases with the increase in solute concentration. It also represents that there is weak interaction between the solute and solvent molecules [17].

The relaxation time (τ) increases with increase in mole fraction, which is in the order of 10⁻¹² sec., is due to structural relaxation process [18] and in such a situation, it is suggested that, the molecule gets rearranged due to co-operative process [19]. The Gibb's free energy (ΔG) increases with increase in mole fraction of the solute, which may due to intermediate compound formation between the ternary liquid mixtures.

As benzene is non- polar molecule doesn't possess dipole moment, when it interacts with chlorobenzene which is polar molecule possess dipole moment then benzene posses induced dipole moment. This is the induced dipole - dipole interaction between benzene and chlorobenzene molecule. Also there is dipole – dipole interaction between the chlorobenzene and Tetrahydrofuran molecule. Hence there exists dipole-dipole interaction between the chlorobenzene and Tetrahydrofuran molecule and dipole – induced dipole interaction between chlorobenzene-benzene molecules.

CONCLUSION

From ultrasonic velocity, related acoustic parameters for ternary mixtures of chlorobenzene with Tetrahydrofuran in benzene for various concentration at 303 K, it has been found that there exists a dipole- induced dipole interaction between chlorobenzene and benzene while dipole-dipole interaction between the chlorobenzene and Tetrahydrofuran molecule.

Acknowledgement

The authors are thankful to Head of Department of Physics, Anand Niketan College, and Warora for providing necessary facility.

REFERENCES

- [1] Kannappam V and Chidambara Vinayagam S, Indian J. Pure & Applied Physics, 2006, 44,670-676.
- [2] Voleisiene B and Voleisis A, Ultragarsas (Ultrasound), 2008, 63(4), 7-9.
- [3] Jain DVS and Dhar N S, Ind. J. Tech, 1992, 3(10), 620.
- [4] Large Man R J and Dunbar W S, J. Phys. Chem., 1945, 49, 428.

[5] Bhandakkar V D, *IOSR Journal of applied Physics*, **2012** 1(5), 38-43.

[6] Bedare G R, Bhandakkar V D and Suryavanshi B.M, International Symposium, Material science and Engineering (IOP science journal), 2012, 42,012028.

[7] Bedare G R, Bhandakkar V D and Suryavanshi B.M, Der Chemica sinica, 2013,4(1), 132-136.

[8] Bedare G R, Bhandakkar V D and Suryavanshi B.M, Der Pharmacia sinica, 2013,4(1), 97-101.

[9] Mistry A A, Bhandakkar V D and Chimankar O P, J. of Chemical & Pharmaceutical Research, **2012** 4(1), 170-174.

[10] Bhandakkar V D, Chimankar O P and Mistry A A, Pelagia Research Library, 2011, 2(6), 70-76.

[11] Pal A & Das G, J. Pure & Applied Ultrasonic, 1990, 21, 9.

- [12] Tabhane V A & Patki B A, Indian J. Pure & Applied Physics, 1985, 23, 58.
- [13] Bhandakkar V D, Adv. Applied Sci. Res, 2011, 2(3), 198-207.
- [14] George J, Sastry N V, Patel S R, &Valand M K, J. Chem. Eng, 2002, 47, 262.
- [15] Bhandakkar V D, Tabhane V A & Ghosh S, Indian J. Pure & Applied Physics, 2003, 41, 849-854.
- [16] Bhandakkar V D, Chimankar O P & Power N R, J. of Chemical & Pharmaceutical Res, 2010, 4, 873-877.
- [17] Lin W and Tsay S J, J. Phys. Chem, 1990, 74, 1037.
- [18] Kinsler LE & Rray A R, Fundamental of Acoustic Wiley eastern, New Delhi, 1989.
- [19] Hyder A and Nair A K, Indian journal of pure and applied Physics, 2000,63,7413