

Total protein and lipid content in edible tissues of fishes from Kasimodu fish landing centre, Chennai, Tamilnadu

***K. Sujatha¹, A. Anitha Joice² and P. Senthilkumaar³**

¹*Dr. M G R Educational and Research Institute University, Chennai*

²*School of Enzymology and Environmental Toxicology, Chennai*

³*Sir Theagaraya College, Chennai, Tamilnadu*

ABSTRACT

*The study focuses on total protein and total lipids – cholesterol, High Density Lipoprotein (HDL), Very Low Density Lipoprotein (VLDL), Low Density Lipoprotein (LDL) and Triglycerides (TGL) in ten fishes from Kasimodu fish landing centre. The edible tissues; muscle and brain were subjected for investigation. The muscle of *Rastrelliger kanagaruta* and the brain of *Arius caelatus* found to be rich in protein content. The highest cholesterol value was recorded in muscle of *Spyraena obtusata*, and in the brain of *Arius caelatus*. The lowest cholesterol was recorded in the muscle of *R. kanagaruta* and in the brain of *Saurida tumbil*. The muscle of *Nemipterus japonicus* and the brain of *A. caelatus* revealed high HDL and it was low in the muscle of *Upeneus sulphureus* and in the brain of *S. tumbil*. More VLDL was observed in the muscle of *Sardinella longiceps* and in the brain of *R. kanagaruta* and low VLDL was observed in the muscle of *N. japonicus* and in the brain of *S. obtusata*. The muscle of *A. caelatus* and the brain of *Pampus niger* were rich in LDL and very meagre LDL was observed in muscle of *P. niger* and in the brain of *S. obtusata*. The muscle of *S. longiceps* and the brain of *R. kanagaruta* showed more amount of TGL and the muscle of *N. japonicus* and the brain of *P. niger* showed the least amount of TGL. The results emphasize the importance of food value among the fishes of Kasimodu fish landing centre and it alarms the consumer a better awareness about the nutritive value of fishes in selection process of edibility and health concern.*

Key words: Protein, Lipid, Nutritive value.

INTRODUCTION

The advent of Blue Revolution has become one of the man's great hopes for future food supplies as the human population multiplies and industrialization increases the problem of environmental pollution. Fishes often referred to as "rich food for poor people" provides essential nourishment especially proteins of high biological values and fat.

Fish is highly nutritious, tasty and easily digestive. It is much sought after by a broad cross-section of the world's population, particularly in developing countries. It is estimated that around 60 percent of people in many developing countries depend on fish for over 30 percent of their animal protein supplies. However, with the increased awareness of the health benefits of eating fish and then ensuing rise in fish species these figures are rapidly changing. Fish also contains significant amounts of all amino acids particularly lysine. Fish protein can be used therefore to complement the amino acid pattern and improve the overall protein quality of mixed diet [1]. In recent years the nutritional importance of aquatic food has increased substantially because of scientifically recognized beneficial effects of eating aquatic food, fats and oil. Fish contributes enormously to the supply of both macro and micro nutrients in our diet [2].

Malnutrition is a general term that indicates a lack of some or all nutritional elements necessary for human health. Protein Energy Malnutrition- the lack of enough protein and food that provides energy which all the basic food groups provide is the most lethal form of malnutrition/ hunger. It is basically a lack of calories and protein.

Fish protein contains all the essential amino acids in required proportions and hence has a high nutritional value, which contribute to their high biological value. Fish is highly proteinous food consumed by the populace. A larger percentage of consumers do eat fish because of its availability, flavoring and palatability while fewer do so because of its nutritive value [3].

Lipids and their components play vital role in the biochemical adaptations of living organism that dwell in the severe and unique conditions of the Northern latitudes. These lipids are of major importance in the ecological and biochemical monitoring and testing of aquatic organisms. The role of lipids in cellular metabolism is versatile, although three main functions have been identified: energetic; structural and bioeffector roles (i.e. lipids acting as messengers) [4]. Fish lipid contains long-chain n3 (omega-3) PUFA, particularly EPA and DHA. Consumptions of these PUFA's have been perceived to be important in human nutrition, health and disease prevention. World fish lipid request continue to increase. Cholesterol is undoubtedly the most publicized lipid in nature, because of the strong correlation between high levels of cholesterol in seasonal differences and significant losses may occur during processing and storage of foods [5].

MATERIALS AND METHODS

Ten fishes namely *Anchoviella indica*, *Arius caelatus*, *Mugil cephalus*, *Nemipterus japonicus*, *Pampus niger*, *Rastrelliger kanagurta*, *Sardinella longiceps*, *Saurida tumbil*, *Spyraena obtusata* and *Upeneus sulphureus* were selected from East Coast of Northern Tamilnadu, Kasimodu fish landing centre. Total Protein, Total Lipid – Total Cholesterol, High Density Lipoprotein (HDL), Very Low Density Lipoprotein (VLDL), Low Density Lipoprotein (LDL,) and Triglycerides (TGL) contents were studied.


Brain and Muscle were taken from all the fishes selected and approximately 100 mg of tissues were weighed, homogenized in the saline medium and centrifuged. The supernatant was taken for further studies. The protein content was estimated by the dye binding method of [6], Total Cholesterol [7], Estimation of Lipoproteins [8], Estimation of Triglycerides [9], Estimation of HDL [10].

RESULTS AND DISCUSSION

Total Protein content of the fishes namely *A. indica*, *A. caelatus*, *M. cephalus*, *N. japonicus*, *P. niger*, *R. kanagurta*, *S. longiceps*, *S. tumbil*, *S. obtusata* and *U. sulphureus* were represented in Table:1 and Figure: 1. Total Cholesterol content in muscle and brain of ten fishes were picturized in Table: 2. The data of High Density Lipoprotein level in muscle and brain of ten fishes is furnished in Table:3. Very Low Density Lipoprotein values of both the tissues are tabulated in Table 4 and Low Density Lipoprotein values of both the tissues are tabulated in Table 5. Information about Triglycerides in both muscle and brain tissues of ten fishes from Kasimodu fish landing centre is shown in Table:6. The variations among the fishes with reference to different lipid parameters were represented in Figure: 2.

Table 1: Total protein content ($\mu\text{g}/100\text{ mg}$ of tissue)

FISHES	MUSCLE	BRAIN
<i>A. indica</i>	1118.27 \pm 1.40	14.07 \pm 3.04
<i>A. caelatus</i>	138.07 \pm 1.72	38.60 \pm 1.11
<i>M. cephalus</i>	139.47 \pm 0.72	20.47 \pm 1.30
<i>N. japonicus</i>	142.27 \pm 1.72	18.27 \pm 1.50
<i>P. niger</i>	116.40 \pm 5.10	13.53 \pm 0.95
<i>R. kanagurta</i>	180.07 \pm 1.89	24.93 \pm 1.72
<i>S. longiceps</i>	67.50 \pm 1.11	25.93 \pm 3.19
<i>S. tumbil</i>	99.47 \pm 0.70	24.6 \pm 1.94
<i>S. obtusata</i>	113.60 \pm 0.80	17.47 \pm 0.83
<i>U. sulphureus</i>	157.27 \pm 1.47	12.13 \pm 1.70

Figure 1: Total protein content ($\mu\text{g}/100$ mg of tissue)Table 2: Total cholesterol content ($\mu\text{g}/100$ mg of tissue)

FISHES	MUSCLE	BRAIN
<i>A. indica</i>	22.26 \pm 0.41	41.90 \pm 1.47
<i>A. caelatus</i>	26.80 \pm 0.72	86.40 \pm 1.31
<i>M. cephalus</i>	26.26 \pm 0.90	51.03 \pm 1.45
<i>N. japonicus</i>	22.86 \pm 1.30	53.20 \pm 0.80
<i>P. niger</i>	25.65 \pm 0.55	69.73 \pm 0.70
<i>R. kanagurta</i>	20.16 \pm 0.99	72.00 \pm 1.70
<i>S. longiceps</i>	27.20 \pm 0.87	53.53 \pm 1.13
<i>S. tumbil</i>	24.86 \pm 0.70	41.20 \pm 0.87
<i>S. obtusata</i>	27.20 \pm 0.81	54.83 \pm 0.77
<i>U. sulphureus</i>	20.53 \pm 0.70	41.56 \pm 1.76

Table 3: High Density Lipoprotein ($\mu\text{g}/100$ mg of tissue)

FISHES	MUSCLE	BRAIN
<i>A. indica</i>	12.12 \pm 1.37	28.22 \pm 1.66
<i>A. caelatus</i>	12.56 \pm 1.47	45.24 \pm 3.40
<i>M. cephalus</i>	1.73 \pm 1.22	29.53 \pm 1.07
<i>N. japonicus</i>	15.13 \pm 1.11	38.35 \pm 1.71
<i>P. niger</i>	13.24 \pm 1.24	38.27 \pm 1.19
<i>R. kanagurta</i>	11.71 \pm 1.86	38.04 \pm 1.74
<i>S. longiceps</i>	10.28 \pm 1.17	34.47 \pm 2.41
<i>S. tumbil</i>	12.37 \pm 1.21	28.85 \pm 1.18
<i>S. obtusata</i>	14.53 \pm 0.67	40.66 \pm 1.26
<i>U. sulphureus</i>	7.39 \pm 1.24	28.20 \pm 1.97

Table 4: Very Low Density Lipoprotein ($\mu\text{g}/100$ mg of tissue)


FISHES	MUSCLE	BRAIN
<i>A. indica</i>	10.43 \pm 0.90	11.13 \pm 0.27
<i>A. caelatus</i>	8.49 \pm 0.71	13.04 \pm 0.20
<i>M. cephalus</i>	9.19 \pm 0.95	12.01 \pm 0.15
<i>N. japonicus</i>	7.12 \pm 0.98	11.45 \pm 0.85
<i>P. niger</i>	10.11 \pm 0.35	11.51 \pm 0.63
<i>R. kanagurta</i>	8.05 \pm 1.31	16.20 \pm 0.49
<i>S. longiceps</i>	12.56 \pm 0.93	14.30 \pm 0.62
<i>S. tumbil</i>	8.75 \pm 0.31	14.62 \pm 0.60
<i>S. obtusata</i>	10.23 \pm 0.18	10.47 \pm 1.08
<i>U. sulphureus</i>	7.72 \pm 0.62	11.14 \pm 1.50

Table 5: Low Density Lipoprotein ($\mu\text{g}/100$ mg of tissue)

FISHES	MUSCLE	BRAIN
<i>A. indica</i>	0.99 \pm 0.34	2.52 \pm 0.80
<i>A. caelatus</i>	5.73 \pm 0.80	14.49 \pm 1.32
<i>M. cephalus</i>	4.29 \pm 0.65	7.96 \pm 1.85
<i>N. japonicus</i>	1.22 \pm 0.37	4.64 \pm 0.86
<i>P. niger</i>	0.93 \pm 0.41	19.86 \pm 1.26
<i>R. kanagurta</i>	0.99 \pm 0.24	16.49 \pm 1.24
<i>S. longiceps</i>	1.46 \pm 0.60	4.45 \pm 1.05
<i>S. tumbil</i>	2.38 \pm 0.59	1.91 \pm 1.03
<i>S. obtusata</i>	1.16 \pm 0.23	1.56 \pm 0.47

Table 6: Triglycerides ($\mu\text{g}/100$ mg of tissue)

FISHES	MUSCLE	BRAIN
<i>A. indica</i>	69.29 \pm 0.73	54.86 \pm 0.30
<i>A. caelatus</i>	63.32 \pm 1.13	65.77 \pm 0.33
<i>M. cephalus</i>	59.13 \pm 1.76	59.63 \pm 0.86
<i>N. japonicus</i>	58.22 \pm 1.55	59.63 \pm 0.86
<i>P. niger</i>	61.10 \pm 0.96	50.58 \pm 0.68
<i>R. kanagurta</i>	67.33 \pm 1.42	79.50 \pm 1.57
<i>S. longiceps</i>	98.55 \pm 1.59	74.43 \pm 0.77
<i>S. tumbil</i>	69.54 \pm 1.04	79.36 \pm 1.02
<i>S. obtusata</i>	75.11 \pm 1.18	53.36 \pm 1.32
<i>U. sulphureus</i>	60.47 \pm 1.01	59.84 \pm 1.16

Figure 2: Variations in lipid content in muscle and brain of ten fishes from Kasimodu fish landing centre ($\mu\text{g}/100$ mg of tissue)

Biochemical studies are very important from the nutritional point of view. Protein is essential for the sustenance of life and accordingly exists in the largest quantity of all nutrients as a component of the human body [11]. In various fish species, proteins are of important as structural compounds, biocatalysts and hormones for control of growth and differentiations [12]. Protein in fish is a main component constituent of tissue and organs. They are precursors of other nitrogen compounds (enzymes, hormones, slurry, neurotransmitters, cofactors, etc) and constitute an important energy source. The effect of dietary lipid levels on fish growth performance varies considerably within species, size, age, diet and composition, range of lipids level tested and rearing conditions [13]. Inadequate protein levels in the diets result in a reduction of growth and loss of weight. However, when an excess of protein is supplied in the diet, only part of it is used for protein synthesis (growth) and the remaining is transformed into energy [14].

Each body cell is composed mainly of protein. Protein makes up the membrane surrounding the cell and also occurs within the cell. During growth period, adolescence and pregnancy, the number of cell increases and more protein is required for cell growth. In all stages of life tissue protein is constantly being broken down and must be replaced by dietary protein. Protein plays a vital role in the formation of enzymes, antibodies and hormones and other substances that regulate the body process.

Fish and shellfish are important source of protein and income for people in Southeastern Asia [15]. They are also increasingly marketed for the health benefits to consumers [16]. The requirement of nitrogen and sulphur is regulated by dietary protein. The protein immunoglobins act as prime defense against bacterial and viral infections. Proteins by means of exerting osmotic pressure help in maintenance of electrolyte and water balance in human system. Several studies show that protein derived from fish, balances many body regulatory factors.

It is well known that protein is the most important and expensive item that should be supplied in adequate amounts to support good growth with minimal cost [17], [18]. [19] determined the proximate composition and energetic values of selected marine fish and shellfish from West Coast of Peninsular Malaysia. This study has included 20 species of fish 10 pelagic fish and 10 demersal fish. The study revealed that Long tail shad (Terunk) contained high fat content. Long-tailed butterfly ray contained the highest protein. Similarly in our study the fish *Rastrelliger kanagurta* is found to be a healthy choice because it contains highest amount of protein and lowest quantity of cholesterol. The brain of *Arius caelatus* is found to be effective because it contains high protein content, high cholesterol value and high HDL. According to the work done by [20] in Catfish it is clear that antimicrobial proteins and peptides play key role in innate immunity and they had been observed from a wide variety of organisms in last few years. Hence, the fishes rich in protein will produce more innate immunity. *R. kanagurta* would be more useful in developing innate immunity.

Cholesterol is undoubtedly the most publicized lipid in nature, because of the strong correlation between high levels of cholesterol in the blood and the incidence of disease of the cardiovascular system in humans. Usually, the cholesterol content will be more in fish liver oils but in the present investigation the consumable part of fish, muscle and brain were found to contain cholesterol. It is the essential constituent of cells. It aids in the permeability of the cells. It controls the red cells from being easily homolyzed. It functions as the defensive action and transports fat to liver in the form of cholesterol ester for oxidation. It assists the formation of bile acids and bile salts, 7-dehydrocholesterol and vitamin D3, corticosteroid hormone, androgens, estrogens and progesterone. Cholesterol helps the granulation of cell division and acts as an antagonist to phospholipids.

High Density Lipoprotein (HDL) transports cholesterol and its esters from peripheral tissues to the liver for its catabolism (scavenging action). Very Low Density Lipoprotein (VLDL) transports mainly endogenous triglycerides synthesized in hepatic cells from the liver to the extra-hepatic tissue including adipose tissue for storage. Low Density Lipoprotein (LDL) regulates cholesterol synthesis in extra-hepatic tissue. The triglycerides are the most abundant of all lipids. They constitute about 98% of total dietary lipids, the remaining 2% consists of phospholipids and cholesterol and its ester. They are major components of storage or depot fats in animal cells but not normally found in membranes. Triglycerides can be stored in quantities, sufficient to supply the energy needs of the body for many months as in the case of obese person. They are not only stored for longer duration but also yield over twice as much energy as carbohydrates.

Lipids and fatty acids play a significant role in membrane and have a direct impact on membrane mediated process such as osmoregulation, nutrient assimilation and transport. On the other hand, the nature and quantity of these lipids in fish vary according to species and habit [21]. Previous studies correlate with our present investigation pertaining to lipid observations.

CONCLUSION

Nutrition is core pillar of human development. High prevalence of low birth weight, high morbidity and mortality in children and poor maternal nutrition of the mother continue to be major nutritional concerns in India. Clearly, this situation emphasizes the need for examining several issues of nutritional significance. One among the issues is the dearth of information about the nutritional benefits of our food. Fish is mostly consumed for its delicacy with little knowledge of its nutritional wealth. The current examination on the nutritional profile of ten edible and economical fishes provides a substantial range of nutritional information, bringing to our attention the richness of healthy nutrients present in the eatable portion such as muscle and brain. Since it is an attempt on locally available and affordable fishes, a larger section of population of Coromandal coast can reap benefits out of this investigation. The study concludes that locally obtainable fish food can be a substantial aid in redressing the problems of malnutrition in our country.

REFERENCES

- [1] FAO, 2005, United Nations Food and Agriculture Organization, Nutritional elements of fish, FAO, Rome.
- [2] Flowra AF, Tumpa SA, DAV *International Journal of Science* Volume-1, Issue-2 June, 2012, ISSN, 2277-5363.

- [3] Wahengbam Sarjubala Devi, Ch Sarojnalini, *International Journal of Science and Research (IJSR)*,**2013**, India Online ISSN, 2319-7064, Volume 2 Issue 2.
- [4] Svetlana AM, Zinaida AN, Stig Falk Peteren, Pauli OR, Tatiana RR, Svetlana NP, Nina NN, *International Journal of Molecular Science*,**2013**, 14, 7048-7060, doi, 10.3390/ijms14047048.
- [5] Spinelli J, Dassow JA, In Chemistry and Biochemistry of Marine Food Products, *AVI Publishing Company*, Westport, Connecticut,**1982**, pp13-37.
- [6] Bradford MM, *Annal Biochem*,**1976**, 72, 248-254.
- [7] Parekh AC, Jung DH, *Anal Chem*,**1970**, 42, 1423-1427.
- [8] Wilson DE, Spiger MJ, *Journal of Laboratory Clinical Med*,**1973** 82, 473-483.
- [9] Rice EC, *Standard Method of Clinical Chemistry*,**1973**, 6, 215-222.
- [10] Fisenberg, Fredrickson, Catel, *Standard method for Practical Biochemistry*,**1984**, 8, 314-319.
- [11] Sudhakar M, Raja K, Anathan G, Sampathkumar P, *Asian Journal of Biological Sciences*,**2011** 4(2), 166-174,ISSN, 1996-3351/ DOI, 10.3923/ajbs.2011.166.174.
- [12] Amal MY, Naheb SG, *International Journal of Environmental Science and Engineering (IJESE)*,**2012**, ISSN, 2156-7549, 2156-7549 Vol.3, 1-10.
- [13] Arredondo Figueroa JL, Matsumoto Soule, Ponce Palafox JL, Shiral Matsumoto, Gomez Marquez JL, *International Journal of Animal and Veterinary Advances*, **2012**, 4(3), 204-2013, ISSN, 2041-2908.
- [14] Diane Kpogue, Herman Gangbazo, Emile Fiogbe, *Turkish Journal of Fisheries and Aquatic Sciences*,**2013**, 13, 111-117, DOI, 10.4194/1303-2712-v13_1_14.
- [15] Agusa T, Kunito T, Sudaryanto A, Monirith I, Kan Atrireklpap S, Iwata H, Isamil S, Sanguansin J, Muhtar M, Tana TS, Tanabe S, *Environmental Pollution*,**2007**, 145(3), 766-777.
- [16] Schmidt EB, Rasmussen LH, Rasmussen JG, Joensen AM, Madsen MB, Christen JH, Fish marine acids and coronary heart disease, Prostaglandin, Leukotrienes and Essential Fatty acids,**2006**, 73(3),191-195.
- [17] Wec KL, Tacon AGJ, *Bull Jap Soc Sci Fish*,**1982**, 48, 1463-1468. Doi,10.2 331/ suisan 48, 1463.
- [18] Zehra S, Khan MA, *Aquacult Int*,**2011**, 20(2), 383-395, Doi, 10.1007/s 10499-011-9470-8.
- [19] Nurnadia AA, Azrina A, Amin I, *International Food Research Journal*,**2011**, 18, 137-148.
- [20] Anbuhezhan RJ, Gobinath C, Ravichandran S, *World Applied Sciences Journal*,**2011**, 12(3), 256-260.
- [21] Kumaran R, Ravi V, Gunalan B, Murugan S, Sundramanickam, *Pelagia Research Library, Adv Appl Sci Res*, **2012**, 2015-2019.