



## Understanding Myocardial Localized Necrosis: Uncovering the causes behind Heart's Calm Peril

Fang Lu\*

Department of Cardiovascular Medicine, Kumamoto University, Japan

### INTRODUCTION

Myocardial infarction, commonly known as a heart attack, is a life-threatening medical condition that occurs when there is a sudden blockage of blood flow to a part of the heart. This interruption in blood supply leads to the death of heart muscle cells, resulting in various degrees of heart damage. Myocardial infarction is a significant global health concern, claiming millions of lives every year. To effectively prevent and manage this condition, it is crucial to delve into the underlying causes that contribute to its development. This article aims to provide an in-depth exploration of the multifaceted causes of myocardial infarction. Atherosclerosis, a chronic inflammatory condition, is the primary driver behind the majority of myocardial infarctions. It involves the buildup of fatty deposits called plaques within the walls of arteries, causing them to narrow and harden. These plaques consist of cholesterol, fat, calcium, and other cellular debris. Over time, these deposits can rupture, triggering the formation of blood clots that can completely block blood flow to the heart muscle. High blood pressure, or hypertension, is a significant risk factor for myocardial infarction [1,2]. The force of elevated blood pressure puts undue stress on the arterial walls, promoting the development of atherosclerosis.

### DESCRIPTION

Hypertension can also lead to the weakening of blood vessel walls, making them more prone to rupture and the formation of blood clots. Diabetes mellitus, particularly type 2 diabetes, is closely linked to an increased risk of myocardial infarction. Elevated blood sugar levels in diabetes damage blood vessels and contribute to the formation of atherosclerotic plaques. Additionally, diabetes can impair the body's ability to regulate cholesterol levels, further exacerbating the risk of heart disease. Cigarette smoking is a well-established risk factor for myocar-

dial infarction. The toxic chemicals in tobacco smoke damage the inner lining of blood vessels, promoting inflammation and the buildup of plaque. Smoking also reduces the amount of oxygen that can reach the heart, making it work harder to function properly. Dyslipidemia refers to abnormal levels of lipids, such as cholesterol and triglycerides, in the blood. High levels of Low-Density Lipoprotein (LDL) cholesterol, often termed "bad cholesterol", contribute to the formation of atherosclerotic plaques. Conversely, low levels of High-Density Lipoprotein (HDL) cholesterol, known as "good cholesterol," fail to effectively remove excess cholesterol from arterial walls. Lack of physical activity and a sedentary lifestyle contribute significantly to the development of myocardial infarction [3-5]. Regular exercise helps maintain healthy blood pressure, promotes good cholesterol levels, and aids in weight management. Physical activity also improves overall cardiovascular health by enhancing blood flow and reducing inflammation.

### CONCLUSION

Myocardial infarction is a complex condition with a multitude of contributing factors. Addressing these causes through preventive measures is key to reducing the burden of heart disease worldwide. Individuals can lower their risk by adopting a heart-healthy lifestyle, including a balanced diet, regular exercise, stress management, and avoiding smoking. Additionally, early screening and management of risk factors, along with medical interventions when necessary, can play a pivotal role in preventing the occurrence of myocardial infarction. As research continues to advance our understanding of these causes, greater strides can be made in preventing and managing this silent but potentially deadly threat to heart health.

### ACKNOWLEDGEMENT

None.

<b>Received:</b>	01-May-2023	<b>Manuscript No:</b>	IPIC-23-17434
<b>Editor assigned:</b>	03-May-2023	<b>PreQC No:</b>	IPIC-23-17434 (PQ)
<b>Reviewed:</b>	17-May-2023	<b>QC No:</b>	IPIC-23-17434
<b>Revised:</b>	22-May-2023	<b>Manuscript No:</b>	IPIC-23-17434 (R)
<b>Published:</b>	29-May-2023	<b>DOI:</b>	10.21767/2471-8157.9.5.41

**Corresponding author** Fang Lu, Department of Cardiovascular Medicine, Kumamoto University, Japan, E-mail: fang.l@kumamoto-u.ac.jp

**Citation** Lu F (2023) Understanding Myocardial Localized Necrosis: Uncovering the causes behind Heart's Calm Peril. Interv Cardiol J. 9:41.

**Copyright** © 2023 Lu F. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## CONFLICT OF INTEREST

The author's declared that they have no conflict of interest.

## REFERENCES

1. Cai Z, Liu J, Wang H, Yin D, Song W, et al. (2020) Diffuse coronary artery dilation predicted worse long-term outcomes in patients with coronary artery ectasia. *Int J Cardiol.* 319:20-25.
2. McCrindle BW, Rowley AH, Newburger JW, Burns JC, Bolger AF, et al. (2017) Diagnosis, treatment, and long-term management of kawasaki disease: A scientific statement for health professionals from the American heart association. *Circulation.* 135(17):e927-e999.
3. Xu C, Xu L, Gao Z, Zhao S, Zhang H, et al. (2018) Direct delineation of myocardial infarction without contrast agents using a joint motion feature learning architecture. *Med Image Anal.* 50:82-94.
4. Ordovas KG, Higgins CB (2011) Delayed contrast enhancement on MR images of myocardium: Past, present, future. *Radiology.* 261(2):358-374.
5. Thygesen K, Alpert JS, Jaffe AS, Chaitman BR, Bax JJ, et al. (2018) Fourth universal definition of myocardial infarction. *J Am Coll Cardiol.* 72(18):2231-2264.