# iMedPub Journals www.imedpub.com Journal of Healthcare Communications ISSN 2472-1654

2017

Vol. 2 No. 4: 65

DOI: 10.4172/2472-1654.100106

# Using Social Media to Reduce Health Impacts from Extreme Weathers: A Feasibility Study

#### **Abstract**

**Introduction:** Extreme weathers due to climate change have exerted negative impacts on human health. Innovative knowledge dissemination strategies are necessary to reduce the negative impacts of extreme weathers.

**Methods:** We examined the feasibility of social media as a tool to disseminate scientific knowledge regarding extreme weathers to common people. For this purpose, we set up a Facebook (FB) community page, and analyzed various metrics to demonstrate its feasibility as a communication tool that can reach a large number of people in society rapidly.

**Results:** The FB community page was able to reach more than 12,000 viewers within a week. 78% of these viewers were females, and 63% of them aged between 18 and 24. Within New York State (NYS), we have reached viewers from more than 40 cities. Interestingly, we have also reached viewers from other states, and even countries beyond United States.

**Discussion:** Our results demonstrate that social media is a feasible technology to disseminate extreme weathers-related knowledge. However, the technology needs to be used properly for different social groups as target audience.

Keywords: Social media; Public health; Facebook; Twitter

**Received:** September 13, 2017; **Accepted:** September 25, 2017; **Published:** October 05, 2017

#### Introduction

In the face of climate change, people need to be adequately informed of the latest scientific knowledge to reduce the negative health impacts of extreme weathers [1]. Given the pervasiveness of social media in society, the technology may be utilized to disseminate knowledge related to extreme weathers [2]. This project aims to demonstrate the feasibility of using social media to disseminate extreme weathers-related knowledge.

Literature: Researchers have utilized social media for different public health purposes [3]. However, it remains unclear how social media may be used effectively to disseminate knowledge with respect to extreme weathers and climate change [4]. A feasible strategy is to translate academic publications into short messages in laymen language, coupled with pictures and/or graphical illustrations, and disseminate them in popular social media platforms such as Facebook (FB) and Twitter [5].

#### Methods

We developed a FB Community page, and tested the "Promote

## Ricky Leung<sup>1\*</sup>, Shao Lin<sup>2</sup> and Huibin Guo<sup>3</sup>

- Department of Health Policy, Management and Behavior, School of Public Health, University at Albany, SUNY, USA
- 2 Department of Environmental Health Sciences, School of Public Health, University at Albany, SUNY, USA
- 3 School of Economics and Management, Hebei University of Economics and Business, USA

#### \*Corresponding author:

Ricky Leung

rleung@albany.edu

PhD, Assistant Professor, Department of Health Policy, Management and Behavior, School of Public Health, University at Albany, #18, Rensselaer, NY 12144, USA.

Tel: 518-402-6512

**Citation:** Leung R, Lin S, Guo H (2017) Using Social Media to Reduce Health Impacts from Extreme Weathers: A Feasibility Study. J Healthc Commun. Vol. 2 No. 4:65.

Page" and "Boost Post" functions to push selected contents to FB users. Our target audience was residents in New York State (NYS). The posted contents were short messages developed by researchers at the New York State Department of Health (NYSDOH). These messages are scientifically valid, and culturally appropriate for general audience.

We examined the geographical coverage of engaged NY residents, and evaluate the effectiveness of boosting messages in comparison with "organic" dissemination (i.e. open the platform without deliberating pushing these messages). Based on the literature, we also tested the effects of including a picture.

#### Results

The FB community page reached more than 12,000 viewers within 1 week. Among them, 78% were females and 22% were

males. 63% of them aged between 18 and 24. The rest of them were from other age groups.

We focused on the New York State (NYS) for the dissemination effort. Within NYS, viewers came from more than 40 cities. Viewers came from outside NYS as well. Interestingly, some viewers were from other countries, including more than 20 countries outside United States.

Thus, while using FB's paid post boosting was very effective to reach a larger number of viewers, the posted contents could reach viewers "organically" (without being pushed by FB). Other than simply viewing the posted contents, some of these viewers had shown increased engagement by liking, reposting and/or commenting the posted contents.

To increase engagement, the use of well-designed pictures and videos was very effective.

#### **Discussion**

Our results suggest that social media is a useful platform to disseminate extreme weathers-related knowledge and engage the public. More specifically, our FB page reached more female than male users, and more younger than older users. If planning similar projects in the future, these results may serve as very useful building blocks for targeting different social groups. Based on the experience with this project, dedicated staffs and researchers are essential to ensuring that the knowledge and scientific information contained in these platforms are accurate.

#### Limitations

Although our FB community page was able to reach a large number of users, it was unable to show if there was any corresponding behavioral change among these users. Also, we could not ascertain what proportion of the users reached belonged to each specific group of interest. This adds a difficulty to formulate additional strategies to engage social media users to learn about extreme weathers.

#### **Further Research**

An intriguing follow-up study is to examine how different social media platforms are able to engage targeted users for specific purposes, such as increasing emergency preparedness [6]. It is possible that other social media platforms can reach more male and older users [7-10]. We are also interested in studying the effectiveness of different presentation modalities (e.g. pictures, videos, texts) for different knowledge dissemination purposes. Finally, in further research, we seek to collaborate with researchers from other countries suffering from extreme weathers (e.g. China) to tackle this significant public health topic.

## Acknowledgements

We would like to thank the New York State Department of Health (NYSDOH) to provide data for this research.

## **Funding Statement**

We received funding from the Centers for Disease Control and Prevention (CDC) through NYSDOH for this research.

#### References

- 1 Allen KM (2006) Community-based disaster preparedness and climate adaptation: local capacity-building in the Philippines. Disasters 30: 81-101.
- 2 Leung RC (2012) Health information technology and dynamic capabilities. Health Care Manage Rev 37: 43-53.
- 3 Leung R, Li J (2015) Using social media to address Asian immigrants' mental health needs: a systematic literature review. J Nature and Sci 1: 66
- 4 Leiserowitz A, Maibach EW, Roser-Renouf C, Feinberg G, Howe P (2013) Extreme weather and climate change in the American mind.
- 5 Kaplan AM, Haenlein M (2010) Users of the world, unite! the challenges and opportunities of social media. Business Horizons 53: 59-68.
- 6 Keim ME, Noji E (2011) Emergent use of social media: a new age of opportunity for disaster resilience. Am J Disaster Med 6: 47-54.
- 7 Ventola CL (2014) Social media and health care professionals: benefits, risks, and best practices. P T 39: 491-499.
- 8 Denecke K, Bamidis P, Bond C, Gabarron E, Househ M, et al. (2015) Ethical issues of social media usage in healthcare. Yearb Med Inform 10: 137–147.
- 9 Eytan T, Benabio J, Golla V, Parikh R, Stein S (2011) Social media and the health system. Perm J Winter 15: 71–74.
- 10 Snyman L, Visser JH (2014) The adoption of social media and social media marketing by dentists in South Africa. SADJ 69: 260-264.