

March 15-16 2018
Barcelona, Spain

Rajas M Rao et al., Biochem Mol Biol J, Volume 4
DOI: 10.21767/2471-8084-C1-008

DYNAMICS OF TRANSLOCATOR PROTEIN

Rajas M Rao¹, Aurore Vaitinadapoule², Frederic Cadet² and Catherine Ethcebest¹

¹Université de La Réunion,INTS, France

²Universite Paris Diderot, France

Translocator protein (TSPO) is a transmembrane (TM) protein localized in the outer mitochondrial membrane. It consists of five transmembrane helices connected by loops of various lengths. TSPO has been implicated in various pathological situations that include cancer, Alzheimer's and Parkinson's diseases, malaria, inflammation, etc. Initial studies indicated that TSPO would be involved in transport of cholesterol from cytoplasm into the inner membrane of mitochondria. Furthermore, a cholesterol binding site, the so-called CRAC motif was identified. Yet, recent studies have brought contradictory results and led to revisit the actual role of TSPO. Hence, its physiological role remains still unclear. Recent high-resolution structures of TSPO solved for different species in different conditions exhibit significant differences, which confirm that TSPO is a dynamical protein. In the present study, we examine the structural dynamics of the mouse TSPO (mTSPO) protein by means of coarse-grained and all-atom molecular dynamics simulations. As the mTSPO 3D structure was only solved in the presence of the PK-11195 ligand, we address the following questions: i. What are the dynamics and the structure of the protein in the absence of the ligand; ii. What is the impact of the ligand on the dynamics of the protein? Our results show that in absence of ligand TSPO is a highly dynamic protein, characterized by secondary structure deformations. These results

are in very good agreement with solid-state NMR data obtained in the absence of the ligand. In the presence of the ligand, the secondary structures are more preserved, which confirm the stabilizing effect of the ligand. However, surprisingly, the protein exhibits larger fluctuations in the loop regions when the ligand is present. Importantly, we also identified correlated motions between TM helices that differ when TSPO is bound or not to PK-11195. Notably, this ultimately influences the dynamics of the cholesterol-binding CRAC motif. We further propose that these changes in dynamics would have impact on binding properties of TSPO, e.g. binding to cholesterol, or to other proteins that are believed to interact with TSPO as part of various physiological processes.

Biography

Rajas M Rao completed his Master's in Biological Sciences from Bangalore University, India. He worked at Professor R Sowdhamini's laboratory in National Centre for Biological Sciences, Bangalore, on phylogeny of a protein involved in bacterial quorum sensing, LuxS. He is currently working towards his PhD Degree at Inserm unit UMR-1134, on the topic of dynamics and interactions of translocator protein, under the guidance of Professor Catherine Etchebest and Professor Frederic Cadet.

rajasmrao@gmail.com