

February 25-26, 2019
Prague, Czech RepublicYensheng Wang et al., Clin Pediatr Dermatol 2019, Volume: 5
DOI: 10.21767/2472-0143-C1-009

SLIGHT IMPROVEMENT OF THE EFFECTIVENESS IN TREATING CLEFT LIP SCARS WITH SILICONE GEL VERSUS SILICONE SHEET: A SECONDARY ANALYSIS

Yensheng Wang^{1,2} and Qiaowan Lee³¹Yale School of Public Health, USA²Chang-gung Memorial Hospital, ROC³Taipei City Hospital, Taiwan

Introduction: Silicone sheet is often used for preventing cleft lip scar postoperatively with concerns of ingestion, application site irritation and short application time. Silicone gel has day-long application duration with better safety. Previous study suggested no difference between forms of silicone in treating cleft lip scarring without controlling potential confounders. We aimed to evaluate the effectiveness of silicone gel versus sheet in a different model, controlling surgical gap, side and patient effects.

Methods: Data was retrieved from previous clinical trial with additional information from the author. In short, silicone gel/sheets were randomly assigned to the patients. Clinical scar evaluation scores (VSS, VAS, and width) were measured six month after application. Transformation was performed for normalization and direction purpose. Generalized linear model was performed controlling the measurement, age, gap and patient effects with Tukey's adjustment. Sensitivity analysis was performed to compare the result.

Results: There were 29 observations in sheet group with mean VSS 0.46 (SD 0.11), mean 10/VAS 1.42 (SD 1.20), and mean width 2.75 (SD 1.44). There were 26 observations in gel group, with mean VSS 0.41 (SD 0.13), mean 10/VAS 1.35 (SD 0.18), and mean width 2.06 (SD 1.71). Silicone gel had a lower score compared to sheet after Tukey's adjustment ($p=0.0048$, 95 CI: 0.05, 0.26). Sensitivity analysis showed a similar result.

Conclusion: Silicone gel appeared to slightly improve post-operative cleft lip scars compared to silicon sheet. With the advantage of safety and patient-friendly features, silicone gel could be recommended for postoperative care of cleft lip scars in infants.

Biography

Yensheng Wang has completed his MD at Chang-gung Memorial Hospital, Taiwan in 2015. He devoted to clinical studies that discover potential treatments for surgical wounds. He also engaged in discovering investigation techniques and treatment for allergic diseases and cancers. He is current a Graduate student at Yale School of Public Health.

cliff25433@hotmail.com