

July 05-07, 2018  
Berlin, GermanyArch Chem Res 2018, Volume 2  
DOI: 10.21767/2572-4657-C3-009

## CZE AND HPTLC METHODS FOR SIMULTANEOUS DETERMINATION OF CANDESARTAN IN BINARY MIXTURES WITH AMLODIPINE AND HYDROCHLOROTHIAZIDE

**Hytham Ahmed**

Minoufia University, Egypt

**T**wo methods were developed for the simultaneous determination of the antihypertensive drugs candesartan (CAN), Amlodipine (AML) and hydrochlorothiazide (HCT) in their combined tablet dosage form. Method I: capillary zone electrophoresis (CZE) was performed. Electrophoretic conditions were optimized to improve separation, sensitivity, and rapidity. The proposed method used a fused silica capillary (70 cm × 75 µm id), and the background electrolyte was 40 mM phosphate buffer pH 8, with 9 s injection time. The applied voltage was 30 kV. The diode array detector (DAD) was set at 210 nm for measurement of AML, and CAN, and 225 nm for HCT. The Three compounds were resolved in less than 7 min with migration times 4.18, 5.36, and 6.74 min for AML, HCT, and CAN respectively. The described method was linear over the range of 5-100, 2.5-100 and 2.5-50 µg/mL for AML, HCT and CAN respectively with correlation coefficients > 0.9993. For method II, HPTLC analysis was carried out on aluminum-backed sheet of silica gel using

chloroform, methanol, and ammonia in the ratio (8 : 2 : 0.2) as mobile phase. Retardation factors were 0.35, 0.5, and 0.75 for HCT, CAN, and AML respectively. Quantification was achieved with UV densitometry at 365, 265 and 274 nm for AML, CAN and HCT respectively. The linearity ranges were 0.1-0.7, 0.05-0.5 and 0.025-0.5 µg/spot for AML, CAN and HCT, respectively with correlation coefficients > 0.9992. The analytical performance of both methods was thoroughly validated according to International Conference on Harmonization (ICH) guidelines with respect to system suitability, linearity, ranges, precision, accuracy, specificity, robustness, detection and quantification limits. The proposed CZE and HPTLC methods were successfully applied to estimation of the drugs in laboratory prepared mixtures and in their combined tablets dosage form. No chromatographic interference was observed from the tablets excipients.

hmaahmed@yahoo.co.uk