

6th World Congress and Expo on **Applied Microbiology**
&
8th Edition of International Conference on **Antibiotics, Antimicrobials & Resistance**
&
12th International Conference on **Allergy & Immunology**
October 21-22, 2019 Rome, Italy

Hospital-acquired infections in Ukraine

Yulian Konechnyi, Andrij Albokrinov and Olena Korniychuk
Danylo Halytsky Lviv National Medical University, Ukraine

Introduction: According to WHO, approximately 15.0% of all hospitalized patients suffer from Hospital-Acquired Infection (HAIs). Data from USA and Europe show 1,3% to 2,03% cases of HAIs in hospitalized patients. In Ukraine -5-7.000 proven cases of HAIs and 500.000 unregistered cases per year. One of the methods to prevent HAIs is control strategy in Intensive care unit (ICU), where last-line antibiotics are currently used.

Objective: To define spectrum of microbial flora and antimicrobial sensitivity of HAIs in some ICU in Lviv city, Ukraine.

Methods: It was investigated 124 clinical materials from 60 patients at the public hospitals in Lviv during July 2018 - May 2019. Were isolated 144 strains of pathogens. HAIs was defined as infection that appeared in patient later than 48 hours after hospital admission. Causative bacterial strains were isolated and identified using bacteriological method (Mueller Hinton Agar, Blood Agar, Saburo Agar, selective & differential media CHROMID*VRE, CHROMID*MRSA SMART, CHROMID*Candida etc. (Biomerieux)) and biochemical tests MIKRO-LA-TEST NEFERMtest24, CANDIDAtest21, STAPHYtest24 etc. (ErbaLacema). Antibiotic sensitivity test was done according to Kirby-Bauer technique.

Results: HAIs structure was as follows: *Staphylococcus lentus* (17.7%), *Escherichia coli* (20.2%), *Klebsiella pneumoniae* (9,7%), *Pseudomonas aeruginosa* (24.2%), *Streptococcus spp.* (2.4%), *MRSA* (3,2%), *Candida albicans* (8.1%). There were several types of HAIs which are rarely founded, such as *Staph.simulans*, *Staphylococcus gallinarum*, *Staphylococcus haemolyticus* *Staphylococcus lugdunensis*, *Aerococcus viridans*, *Pragia fontium*, *Aeromonas ichthiosmia*, *Raoultella terrigena*, *Burkholderia pseudomallei*, *Macrococcus caseolyticus*, *Pseudomonas oryzihabitans*, *Pseudomonas putida*, *Candida dubliniensis*, *Candida membranifaciens*, *Candida glabrata*. Anti-microbial sensitivity assessment showed that more than 90.0 % pathogens were multidrug resistance (MDR). 75% of strains were atypical strains (T-index $\geq 0,25$).

Conclusion: In the era of antibiotics HAIs are still uncontrollable. Most cases of HAIs in Ukraine are not recognized as epidemics. More than 90.0 % pathogens are MDR and most of them are atypical strains. Diagnostic capabilities show a number of rare species of microorganisms that cause nosocomial infections. It is important to use this fact when one attempts to design prevention or control strategy to reduce HAIs either at WHO level, or at the level of local health care unit.

Acknowledgement: We thank directors of the hospitals and ICU teams.

Ethical Committee: 25/06/2018 № 6.

Biography

Yulian Konechnyi is doing his PhD in microbiology at Danylo Halytsky Lviv National Medical University, Ukraine. He has completed his MD and have specialization in internal medicine. He has worked as lecturer at the department of microbiology in microbiology, virology and immunology topics, at Danylo Halytsky Lviv National Medical University, Ukraine. He has published 10 different papers.