

Research Article

Nutritional Deficiency Disorders Associated with Pregnancy

Mehwish Iftikhar*

People Sourcing Specialist, Synavos Lahore District, Punjab, Pakistan

Abstract

Maternal paleness is a typical issue in pregnancy, especially in developing nations. Pallor influences very nearly 66% of pregnant ladies in creating nations and adds to maternal dismalness and mortality and to low birth weight. Pregnant ladies establish a high hazard gather for insufficiency. Micronutrient lack particularly the iron, folate, calcium and zinc inadequacy is the most despicable aspect of our lives, influencing all areas of society. Lamentably, the ladies amid pregnancy, youthfulness, and youngsters are under this revile especially in creating nations like Pakistan. A high level of the pregnant ladies is iron inadequate because of elements, for example, high equality, poor dietary accessibility and financial status. Intercessions to improve the admission of a few micronutrients ought to be considered to counteract iron deficiency in pregnant ladies. The systems and battle of Pakistani government will prompt a positive bring about killing micronutrient lack from Pakistan. Nutritional status assumes vital job in anticipating micronutrient

inadequacy. There is a critical need to expand mindfulness with respect to significance of this less appetite among WOCBA particularly by including medicinal services experts, network wellbeing laborers, drug specialist, media, symposiums furthermore, workshops as it is totally preventable general medical issue. In the 1st trimester the percentages of values falling into the critical range are significantly increased for thiamine and vitamin B6, but not for vitamin A, riboflavin, and vitamin C. In the 3rd trimester an additionally greater number of women not sufficiently supplied could be found for vitamin A; the percentages of values for thiamine and vitamin B6 in the critical range were found to be significantly higher. For iron and folate, likewise known to be critical nutrients in pregnancy, no supply parameters can be presented at the present time from our investigations.

Keywords: Maternal paleness; Nutritional deficiency; Nutritional disorders; Pregnancy

Introduction

In this challenging environment, the number of inhabitants in developing nations is on crest than world's normal populace. There is a need of genuine hard exertion with full sustenance insurgency particularly in poor and creating nations to illuminate nourishment shortage and accessibility [1], yet it requires steady and decided endeavors all things considered and researchers all around the globe. In correlation of absolute populace, about 65% of world's all out inhabitants is starving and lives underneath the neediness line. Hunger and under sustenance are found among individuals where nourishment supply and diet diversification are inadequate. In this 21st century, the creating scene is under extreme danger of micronutrient lack of healthy sustenance in light of the utilization of eating regimen that are rich in phytates and these points of confinement bioavailability of fundamental supplements [2]. The lack of basic nutrients and minerals are currently viewed as intensive and it influences more than 33% of the total populace, effective ladies and youngsters, coming about destroying outcomes for general wellbeing, and social development of the country. These micronutrient insufficiencies contribute fundamentally to the weight of maladies and connected to antagonistic useful results, for example, hindering, squandering, and expanded defenselessness to diseases amid pregnancy, diminished IQ level, subjective misfortunes, visual impairment, and untimely mortality [3]. Frailty complex pregnancy undermines the life of both mother and foetus. Maternal and fetal mortality and dreariness are related with this issue. Sickliness is exceedingly common in the immature world and it's incidents are found in ladies of low financial class

and among fantastic multiparae. Earlier research in Pakistan reports iron inadequacy as the main source of sickliness in pregnancy. Iron deficiency anemia (IDA) is known to be the most regular nourishing lack around the world. Pakistan, with its crucial geopolitical significance, is as yet far off from its objective to beat IDA among helpless populace gatherings. A mind greater part of Pakistani populace experiences IDA [4]. Typically, all pregnant ladies and nursing moms in Pakistan are influenced by IDA. Predominance of IDA among females of regenerative age exists up to half, hence, maternal death rate is high and a developing concern has been raised by the neighborhood and universal associations to control IDA in Pakistan. As per Pakistan Demographic and Health Survey 2006-07, maternal mortality was appeared to be high with 276 passing's for each 100,000 live births when contrasted with 1 out of 8,000 in the created world. Among 89 Pakistani ladies, one kicks the bucket of labor confusions.

Folate is a normally happening nutrient while folic corrosive is the manufactured substitution of folate utilized in many enhancements and in strengthened nourishments. People are completely reliant on dietary sources or dietary enhancements and microorganisms in their intestinal tract for their folate supply [5]. Folate subordinates are fundamental for the combination of nucleic corrosive, amino acids, cell division, tissue development, and DNA methylation. Insufficient folate admission prompts a diminishing in serum folate fixation, bringing about a reduction in erythrocyte (red platelet) folate fixation, an ascent in homocysteine focus, and megaloblastic changes in the bone marrow and different tissues with quickly separating cells during pregnancy, fetal development causes an expansion in the complete number of quickly partitioning

cells, which prompts expanded necessities for folate [6]. With insufficient folic corrosive admission, convergences of folate in maternal serum, plasma, and red platelets decline from the fifth month of pregnancy onwards. If lacking folate admission is continued amid pregnancy, megaloblastic pallor (a blood issue described by frailty, with red platelets that are bigger than typical and cell substance that are not totally created) happens [6,7].

Zinc insufficiency is a regularly commonly medical issue all through the world [8]. Low pay family's youngsters in Pakistan are accounted for to show low zinc level and authorize to low dietary admission of zinc other than other bewildering variables, for example, instruction, financial condition, dietary propensities and poor absorption [9]. All these need legitimate assessment to be considered. As information regarding the matter is alarm especially in the ladies of Pakistan in this age gathering. Subsequently this investigation is intended to evaluate the greatness of the issue of zinc lack as spellbound in the most powerless cross area of our populace for example the ladies of childbearing age (WCBA-Women child bearing age).

Vitamin D lack amid pregnancy has been connected with number of genuine short and long haul medical issues in off spring, including hindered development, skeletal issues, type 1 diabetes, asthma and schizophrenia yet few have investigated the job of maternal vitamin D status and pregnancy results. Vitamin D adequacy is increasingly basic amid pregnancy and lactation [10]. Calcium ingestion increments amid pregnancy because of fetal requests, consequently, expanding the dimension of vitamin D. Pregnant ladies ought to expend multiple times more nutrient D than typical ladies. The new conceived devours its vitamin D stores in the initial two months. On the off chance that vitamin D insufficiency keeps amid lactation, the danger of rickets increments in breastfed newborn children. 25 (OH) 2D levels are expanded from the earliest starting point of pregnancy [11]. Vitamin D is essential for the retention and digestion of calcium and phosphorous.

Iron

Iron plays a focal job in numerous biochemical procedures in the body. The key capacity of iron is to encourage oxygen transport by hemoglobin, the oxygen-carrying pigment of erythrocytes. It is additionally associated with oxygen stockpiling by myoglobin. Iron is imperative for the expansion of all cells including those of the safe framework [12]. Iron inadequacy sickliness (IDA) is known to be the most widely recognized wholesome insufficiency around the world. Iron lack pallor prompts endless mortalities, maternal drain, diminished school execution and diminished efficiency in defenseless populations, 1-4 and is additionally harming amid pregnancy particularly in the nations of immature world [13]. Anemia is profoundly pervasive in the immature World.

Iron deficiency anemia in pregnancy

Pregnancy is a physiological condition and normally has no impact on general health of a pregnant lady. Anyway pregnancy results in hormonal, haemodynamics and hematological changes. These physiological changes should be seen as typical adjustments controlled essentially. Pregnant ladies comprise a high hazard amass for iron lack [14]. Anemia muddling pregnancy undermines the life of both mother and baby. Maternal and fetal mortality and bleakness are related with this issue [15]. A significant assortment

of writing uncovers that IDA is the most predominant dietary insufficiency in Pakistan. Various little examinations demonstrated an extraordinary variety in the degree of pervasiveness of IDA in Pakistan eg 48.2% of the pregnant ladies were appeared to be frail while 90.5% of the absolutetried pregnant ladies experienced IDA. 16 Poor nourishment in rehashed pregnancies and undesirable sustenance propensities have been dominantly connected with the beginning of IDA among helpless populace parts [13]. Earlier research in Pakistan records iron inadequacy as the main source of paleness in pregnancy [15] (Table 1).

Table 1: Cause of anemia in pregnancy.

Symptoms	Number(n=104)	%
Iron deficiency	66	63.5
Folate deficiency	14	13.5
Normocytic normochromic	12	11.5
B12 deficiency	8	7.7
Thalassemia minor	4	3.8

Inadequacy of iron in pregnant ladies limits oxygen bringing to cell bringing resulting fatigue, poor work execution and diminished invulnerability. Iron inadequacy anemia from the get-go in a pregnancy can twofold or even triple the danger of having an unexpected labor or a low birth weight baby. 10When maternal iron stores are exhausted, the embryo can't aggregate much iron and there is a reduction in fetal iron stores. Studies recommend that social anomalies happen in youngsters with iron lack. These variations from the norm are identified with changes in the centralization of synthetic go between in the mind. Iron deficiency in the absence of anaemia is associated with poor performance on Bayley Mental development Index. Advancement delay in iron inadequate babies can be switched by treatment with iron [14].

Dietary modification

Iron insufficiency is the most predominant explicit single miniaturized scale supplement lack influencing around half of the total populace. Among the most influenced by this ailment are pregnant ladies due to included iron necessities amid pregnancy. This is essentially on the grounds that the measure of dietary iron assimilated is frequently too little to even consider meeting the expanded interest amid pregnancy. The predominance of iron insufficiency frailty stays high in numerous pieces of the world in spite of critical exertion to lighten this issue [12]. The national wellbeing study of Pakistan revealed that 43-47% of country and 35-40% of urban ladies matured 15-44 years are iron deficient. The frequency of iron- inadequacy paleness among hopeful moms is alarmingly high. It is a significant issue in pregnancy which influences 50-60% of these ladies. Iron insufficiency is typically the aftereffect of deficient bio accessible dietary iron and expanded iron prerequisites amid a time of quick development in pregnancy and earliest stages. Weakness of pregnancy significantly affectsthe strength of hatchling just as mother, particularly if serious, may hinder the oxygen conveyance to placenta and embryo meddling with intrauterine development. Placental weight, volume and surface zone are decreased if expecting mother is reasonably frail. It results in 12-28% of fetal misfortune, 30% pre-birth passing's and 7- 10% of neonatal passing's. Amid the second trimester sickliness is related with preterm birth, occurrence of which is expanded five- crease for iron lack paleness and twofold for other weakness. The danger of iron insufficiency is especially high in

ladies with high equality and short interims between pregnancies [14].

Iron requirement of human body

The aggregate sum iron lost is assessed to be at 14 µg/kg body weight/day. A non-menstruating 55 kg lady loses about 0.8 mg Fe/day and a 70 kg man loses around 1 mg. The scope of individual variety has been evaluated to be 15% (FAO/WHO). If there should be an occurrence of baby, the necessity of iron is around 75 mg/kg body weight and it just satisfied from human milk that just contain little measure of iron. The prerequisites for ingested iron in babies and kids are high in connection to their vitality necessities. Iron necessities are additionally shifts in youths, especially amid the time of quick development. The mean menstrual iron misfortune, arrived at the midpoint of over the whole menstrual cycle of 28 days, is about 0.56 mg/day [16].

Vitamin D deficiency

Vitamin D inadequacy has been related with rickets, an ailment in which the bone tissue doesn't legitimately mineralize, prompting delicate bones and skeletal disfigurements. Be that as it may, progressively, examine is uncovering the significance of vitamin D in securing against a large group of medical issues. Manifestations of bone agony and muscle shortcoming can mean you have a vitamin D lack [10]. Indeed, even without side effects, too little vitamin D can present wellbeing dangers. Vitamin D insufficiency can happen for various reasons: You don't expend the prescribed dimensions of the nutrient after some time, your introduction to daylight is restricted and you have dull skin.

Vitamin D deficiency during pregnancy

Vitamin D status amid pregnancy is fundamental for the skeletal organization and improvement of baby. Low maternal vitamin D is related with shorter term of development and thusly decreased development of long bones in babies. The essential job of vitamin D is to keep up serum phosphate and calcium levels by advancing their intestinal retention legitimately, or by actuating bone reabsorption through circuitous enlistment and enactment of osteoclasts [11]. Vitamin D adequacy is progressively basic amid pregnancy and lactation. Pregnant ladies ought to expend multiple times more vitamin D than ordinary ladies. The new conceived devours its vitamin D stores in the initial two months. In the event that vitamin D lack keeps amid lactation, the danger of rickets increments in breastfed babies. Amid pregnancy, for the most part in the last trimester, changes in maternal vitamin D and calcium digestion permit the exchange of upto 250 mg of calcium for every day to the fetal skeleton, for an aggregate of 25-30 grams of calcium [11]. Vitamin D lack amid pregnancy has been connected with number of genuine short and long haul medical issues in off spring, including impeded development, skeletal issues, type 1 diabetes, asthma and schizophrenia⁶ yet few have investigated the job of maternal nutrient D status and pregnancy results [10]. Vitamin D insufficiency is related with osteoporosis and an assortment of different sicknesses, running from sadness and extreme myopathy to immune system malady in grown-ups and rickets, newborn child heart disappointment, intense lower respiratory tract infection³ and inappropriate bone advancement at 9 years in youngsters [11].

Dietary modification

It has been assessed that 1 billion individuals worldwide have vitamin D inadequacy or deficiency. Around 70 - 90% of vitamin D lack has been recognized in sound asymptomatic volunteers in two investigations led in Karachi. Studies have demonstrated the commonness of vitamin D lack to be 63.3% in pregnant ladies. Low dimensions of vitamin D amid pregnancy or bosom nourishing can adversely affect infant's development, the arrangement of tooth veneer and the manner in which body handles calcium. It can likewise put an infant in danger of being brought into the world with rickets or creating rickets in adolescence. Newborn children destined to moms with hypovitaminosis D have expanded danger of symptomatic hypocalcaemia, little for gestational age and larger fontanelle, suggestive of impaired ossification of skull bones.

Folate deficiency

Folate is a generic term for both the endogenous form of the vitamin occurring naturally in food and the synthetic form found supplements and fortified foods [17]. It should be noted, however, that folate is a naturally occurring vitamin while folic acid is the synthetic replacement of folate used in most supplements and in fortified foods. Humans are fully dependent on dietary sources or dietary supplements and microorganisms in their intestinal tract for their folate supply. Folate derivatives are essential for the synthesis of nucleic acid, amino acids, cell division, tissue growth, and DNA methylation [18,19]. Inadequate folate intake leads to a decrease in serum folate concentration, resulting in a decrease in erythrocyte (red blood cell) folate concentration, a rise in homocysteine (Hcy) concentration, and megaloblastic changes in the bone marrow and other tissues with rapidly dividing cells [7,20]. During pregnancy, fetal growth causes an increase in the total number of rapidly dividing cells, which leads to increased requirements for folate [17]. With inadequate folic acid intake, concentrations of folate in maternal serum, plasma, and red blood cells decrease from the fifth month of pregnancy onwards [21,22]. If inadequate folate intake is sustained during pregnancy, megaloblastic anaemia (a blood disorder characterised by anaemia, with red blood cells that are larger than normal and cell contents that are not completely developed) occurs [20].

Dietary modification

Subar et al. [23] examined representative data from the second National Health and Nutrition Examination Survey (NHANES II) and found that the estimated mean folate intake of women surveyed (207 ± 2.9 µg/d) was approximately equivalent to the recommended dietary allowance (RDA) for the nonpregnant state (180 µg/d). Approximately 90% of the women consumed <400 µg folate/d (the RDA for pregnancy) and only ≈10% of the women met the pregnancy RDA. More black (26%) than white (18%) women had very low folate intakes (≤ 100 µg/d), potentially accounting for the consistently lower average folate intake reported for minority women (175-185 µg/d). Despite this, there was little or no ethnic difference when daily intake of folate from the diet exceeded 100 µg/d.

Block and Abrams also examined data from NHANES II along with data from the Continuing Survey of Food Intakes by Individuals; they found that women who were near poverty or below poverty had intakes of folate and other nutrients (eg: iron,

zinc, and vitamins A, C, and B-6) that were below the current RDA for non-pregnant women (3 µg folate/kg). One-third of low-income women (<131% of poverty limit) and half of those women with higher incomes (>300% of poverty limit) met the folate RDA for non-pregnant women. Folate intakes below the RDA were associated with infrequent consumption of folate-rich foods. Low-income women in particular ate few vegetables; about half of the women ate no vegetables at all, including potatoes.

The major food sources of folate include cooked dried beans, leafy green vegetables, and fortified cereals [23].

Other foods of lower folate density are also important contributors of folate to the women because such foods are eaten frequently (eg: orange juice and white bread). Multivitamin tablets are an additional source of folate that is used by many people. Orange juice is the largest single source of folate consumed by the public, contributing ≈10% of dietary folate, with white bread, dried beans, salad, and cold cereal contributing a cumulative one-third of folate from diet. One might presume that the use of vitamin and mineral supplements containing folic acid would offset the risk of low folate intake, particularly because the folic acid contained in supplements (eg: monoglutamate) has greater bioavailability than does the folate in food (eg: polyglutamate). However, supplements appear to be used the least by those individuals who need them the most.

Factors associated with improved folate status (measured as red cell folate) include use of fortified cereals in addition to the use of folic acid-containing supplements [24]. The remaining treatments included dietary advice and a control (i.e. no intervention). The intervention treatments significantly raised the folate intake of the women. Even the group receiving dietary advice alone increased their intake by nearly 100 µg folate/d. However, only women taking folic acid supplements or fortified foods had significantly increased red cell folate after 3 months. Likewise, predictors of red cell folate among women attempting pregnancy included use of folic acid-containing nutritional supplements and fortified cereals [25]. Interestingly, there was a substantial ethnic difference in use of folic acid-fortified cereals, ranking 9th as a source of dietary folate among whites, but ranking 49th among black women [23].

Summary

In summary, it suggests that poor dietary folate intake and low circulating concentrations of folate are associated with an increased risk of adverse birth outcomes. Supplementation studies likewise suggest that some women—most likely poor women—may benefit from receiving additional folic acid during, as well as before, pregnancy. Some negative effects have been reported in association with Preconceptional supplementation with folic acid-containing vitamins, including potential increases in spontaneous abortion and infant low birth weight. These risks may be more apparent than they are real, occurring in association with increased fertility, survivorship of marginal conceptions, or a small shift in sex ratio.

Likewise, high concentrations of homocysteine have been associated with increased habitual spontaneous abortion and serious complications of pregnancy, including pregnancy-induced hypertension, preeclampsia, and placental abruption. Currently, there are no data from well-controlled studies to document the

security of these observations or from clinical trials to determine whether supplementation with folic acid and B vitamins reduces the risk associated with maternal hyper-homocysteinemia.

The current WHO guidelines recommend initiation of oral daily IFA supplements as early as possible during the pregnancy (World Health organization Geneva). However, we found that the supplementation initiation was late on average during the fifth month of pregnancy and 17% of women consumed 90 or more supplements throughout their pregnancy. Early initiation and the total number of supplements consumed during pregnancy have a significant impact on child mortality. Hence, early initiation and continued use of supplements is important to reduce neonatal and infant mortality in Pakistan.

Conclusion

Maternal hunger is one of the main sources of various ailments and supplement insufficiencies in youngsters around the globe. The circumstance is most exceedingly bad in the creating nations where there is absence of access to adjusted and broadened diet. Micronutrient lack particularly iron, zinc, folate and calcium insufficiency is a typical ailing health among all. The most powerless gathering of society influenced by micronutrient insufficiency is pregnant ladies and ladies of kid bearing age. The quantities of individuals encountering this micronutrient ailing health are higher in creating nations. Along these lines, the circumstance is very more regrettable in Pakistan. The aversion procedures that can be selected are micronutrient supplementation, sustenance fortress, diet enhancement, and control of malarial and parasitic infections. As indicated by the practicality, any of the apparatuses or blend of these can be executed. The systems and battle of Pakistani government will prompt a positive bring about killing micronutrient lack from Pakistan. Instruction status assumes vital job in anticipating micronutrient inadequacy. There is a critical need to expand mindfulness with respect to significance of this concealed appetite among WOCBA particularly by including medicinal services experts, network wellbeing laborers, drug specialist, media, symposiums furthermore, workshops as it is totally preventable shrouded general medical issue.

References

1. Huang J, Pray C, Rozelle S (2002) Enhancing the crops to feed the poor. *Nature* 418:678-684.
2. Hussain S, Maqsood MA, Rahmatullah A (2010) Increasing grain zinc and yield of wheat for the developing world: A review. *Emir J Food Agric* 22:326-339.
3. Murray-Kolb LE, Beard JL (2009) Iron deficiency and child and maternal health. *Am J Clin Nutr* 89(3): 946S-950S.
4. Lozoff B, Corapci F, Burden MJ, Kaciroti N, Barroso RA, et al. (2007) Pre-school aged children with iron deficiency anemia show altered affect and behavior. *J Nutr* 137:683-689.
5. Hisam A, Rahman MU, Mashhadi SF (2014) Knowledge, attitude and practice regarding folic acid deficiency; A hidden hunger. *Pak J Med Sci* 30(3):583-588.
6. Lassi ZS, Salam RA, Haider BA, Bhutta ZA (2013) Folic acid supplementation during pregnancy for maternal health

- and pregnancy outcomes. *Cochrane Database Syst Rev* 3:CD006896.
7. Dietary Ref 1998 Anonymous (1998) Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic acid, Biotin and Choline. Institute of Medicine, National Academies Press (US), Washington.
 8. De-regil LM, Suchdev PS, Vist GE, Wallester S, Penarosas JP (2011) Home fortification of foods with multiple micronutrient powders for health and nutrition in children under two years of age. *Cochrane Database Syst Rev* 9:CD008959.
 9. Ahmad S, Maqbool S, Mohyidin A (1998) Plasma zinc and copper levels in children. *Pak J Med Res* 27: 148-154.
 10. Ali ST, Naqvi KZ, Maqsood M, Thontia S (2013) Prevalence of Vitamin D Deficiency among Postpartum Women and their Newborns: a cross-sectional study in Karachi, Pakistan. *Pak J Surg* 29(1): 41-45.
 11. Sharif S, Farasat T, Shoaib H, Saqib M, Fazal S (2013) Vitamin d levels among pregnant and lactating women. *J Coll Physicians Surg Pak* 23(12): 862-865.
 12. Al Akhali K, Ali MAH, Ansari MA (2013) Evaluation of prevalence and pattern of anemia—A hospital based study in Aseer Province, Kingdom of Saudi Arabia. *J Exp Med Sur Res* 2: 32-35.
 13. Akhtar S, Ahmed A, Ahmad A, Ali Z, Riaz M, et al. (2013) Iron status of the Pakistani population- current issues and strategies. *Asia Pac J Clin Nutr* 22(3): 340-347.
 14. Raza N, Sarwar I, Munazza B, Ayub M, Suleman M (2011) Assessment of iron deficiency in pregnant women by determining iron status. *J Ayub Med Coll Abbottabad* 23(2):36-40.
 15. Karim SA, Khurshid M, Memon AM, Jafarey SN (1994) Anaemia in pregnancy-its cause in the underprivileged class of Karachi. *J Pak Med Assoc* 44(4):90-92.
 16. Ahmed A, Ahmad A, Khalid N, David A, Sandhu MA, et al. (2014) A question mark on iron deficiency in 185 million people of Pakistan: its outcomes and prevention. *Crit Rev Food Sci Nutr* 54(12):1617-1635.
 17. Bailey LB (1995) Folate requirements and dietary recommendations. In: Bailey LB editor(s). *Folate in Health and Disease*. Marcel Dekker, New York.
 18. Krishnaswam K, Nair KM (2001) Importance of folate in human nutrition. *Br J Nutr* 85: Suppl 2:S115-S124.
 19. Morrison K, Papapetrou C, Hol FA, Mariman EC, Lynch SA, et al. (1998) Susceptibility to spina bifida; an association study of five candidate genes. *Ann Hum Genet* 62(5):379-396.
 20. Willoughby MLN, Jewell FG (1968) Folate status throughout pregnancy and in post partum period. *Br Med J* 4:356-360.
 21. Ackurt F, Wetherilt H, Loker M, Hacibekiro M (1995) Biochemical assessment of nutritional status in pre- and post-natal Turkish women and outcome of pregnancy. *Eur J Clin Nutr* 49(8):613-622.
 22. Bates CJ, Fuller NJ, Prentice AM (1986) Folate status during pregnancy and lactation in a West African rural community. *Human Nutr Clin Nutr* 40:3-13.
 23. Subar AF, Block G, James LD (1998) Folate intake and food sources in the US population. *Am J Clin Nutr* 50: 508-516.
 24. Cuskelly GJ, McNulty H, Scott JM (1996) Effect of increasing dietary folate on red-cell folate: implications for prevention of neural tube defect. *Lancet* 347:657-659.
 25. Brown JE, Jacobs DR, Hartman TJ, Barosso GM, Stang JS, et al. (1997) Predictors of red cell folate level in women attempting pregnancy. *JAMA* 277: 548-552.
- Address of Correspondence:** Mehwish Iftikhar, People Sourcing Specialist, Synavos Lahore District, Punjab, Pakistan, Tel: +03006541006; E-mail: mahwishmalik192@gmail.com
- Submitted: November 08, 2021; Accepted: November 22, 2021; Published: November 29, 2021*