

How Biomarker Indicates Change in Expression or State of A Protein

Zuhal Sezgin*

Department of Medical Biology, Mersin University School of Medicine, Turkey

Abstract

Parkinson's disease (PD) is that the most frequent neurodegenerative movement disorder. In near future its importance can increase, particularly because of changes in age structure. There square measure still substantial issues in terms of early diagnosing of the malady. In clinical observe, the diagnosing of metallic element continues to be supported the analysis of clinical presentation and is supported by varied auxiliary tests with heterogeneous prognosticative price.

Keywords: Parkinson's disease; Pathology; Histopathology

Corresponding author:

Zuhal Sezgin

Tel : +90-122-421-23-01

✉ sezbinzuhal@doc-mum.tr

Received: August 19, 2021; **Accepted:** August 27, 2021; **Published:** September 05, 2021

Citation: Sezgin Z (2021) How Biomarker Indicates Change in Expression or State of A Protein. J Biomark Drug Dev. Vol.2 No.1:102

Introduction

The immune system's illustration of the body. What may be the selective advantage of associate degree system expressing patterns of integral pathology to specific sets of self-molecules? to higher characterize the homunculus, we've used informatic tools to review patterns of associate degreetibodies to several many self-molecules panoplied on glass slides – an substance chip of our style. Halide has been found to cause damaging effects on fish gills [1]. Despite essential roles in varied metabolism activities, whether or not and the way miRNAs participate within the course of toxicity caused by halide in gills continues to be unclear. Triclosan (TCS) may be a broad-spectrum medicament agent that is wide employed in varied care merchandise and cosmetics. It's been found that TCS affects endocrine, immune, nervous, procreative, and organic process system. Though microRNAs (miRNAs) act a crucial half in several metabolic activities, whether or not and the way they're associated with the method of TCS-induced toxicity is unknown [2]. Blubber is that the results of the interaction of multiple variables, as well as the excessive increase of sugar-sweetened beverages consumption. Diets aimed to treat blubber have steered the utilization of artificial sweeteners. However, recent proof has shown many health deficits when intake of artificial sweeteners, as well as effects in neuron activity. Organic phenomenon profiles were organ, duration-, and concentration-dependent, suggesting that stress-response genes may be concerned in associate degree innate arms for handling serious metal exposure [3]. To the most effective of our information, this study is that the initial one in every of histopathology and stress-response organic phenomenon pattern of Scylla paramamosain when Cd exposure. Dexmedetomidine is wide used for perioperative and intensive care unit patients. MicroRNAs (miRNAs) perform as regulators

of organic phenomenon. The aim of the study was to assay expression identification of microRNA in rat hearts following administration of dexmedetomidine [4,5]. The genetic analysis of current tumour cells (CTCs) is going to be applied for preciseness drugs of castration-resistant prostatic adenocarcinoma (CRPC). Previous reports showed that the expression of prostate-specific membrane substance (PSMA) in prostatic adenocarcinoma tissue correlates with cancer aggressiveness. We tend to explore the chance of PSMA expression in CTCs as a prognosticative biomarker for CRPC. Symptom may be a devastating modality disorder impacting a growing range of individuals annually [6]. The aims of this experiment were to assess neuron mechanisms concerned within the initial malleability when traumatic noise exposure that would contribute to the emergence of symptom and to check a possible medical specialty treatment to change this early neural malleability. Inhalation of polyhexamethylene guanidine (PHMG) causes irreversible respiratory organ injury, like respiratory organ pathology. However, the mechanism underlying PHMG-induced respiratory organ injury is unclear.

References

1. Börger M, Funke S, Bähr M, Grus F, Lingor P (2015) Biomarker sources for Parkinson's disease: Time to shed tears. *Basal Ganglia* 5(2-3):63-69.
2. Irun R Cohen (2007) Biomarkers, self-antigens and the immunological homunculus. *J Autoim* 29(4):246-249.
3. Jianjie C, Yongju L, Jinling C, Lingtian X (2021) Fluoride exposure changed the expression of microRNAs in gills of male zebrafish. *Aqua Toxic* 233:105789.

4. Fan W, Fei L, Wanguang C (2019) Exposure to triclosan changes the expression of microRNA in male juvenile zebrafish. *J Autoim* 214:651-658.
5. Mirielde-la-C, Diana MA, Daniela Marcia SN, René DC, Eric MR. (2018) The artificial sweetener Splenda intake promotes changes in expression of c-Fos and NeuN in hypothalamus and hippocampus of rats. *Brain Res* 1700: 181-189.
6. Zhong-Kai Z, Dan-Dan T, Yi-Lian ZW, Ze-Peng L, Wen-Bin G (2018) Effect of cadmium exposure on hepatopancreas and gills of the estuary mud crab (*Scylla paramamosain*): Histopathological changes and expression characterization of stress response genes. *Aqua Toxic* 195:1-7.