

Urinary ¹H-NMR Metabolomics Can Distinguish Pancreatitis Patients from Healthy Controls

Elizabeth R Lusczek¹, Joao A Paulo², John R Saltzman²,
Vivek Kadiyala², Peter A Banks², Greg Beilman¹, Darwin L Conwell²

¹Department of Surgery, University of Minnesota. Minneapolis, MN, USA.

²Center for Pancreatic Disease, Brigham and Women's Hospital, Harvard Medical School. Boston, MA, USA

ABSTRACT

Context The characterization of the urinary metabolome may yield biomarkers indicative of pancreatitis. **Objectives** We establish a non-invasive technique to compare urinary metabolic profiles in patients with acute and chronic pancreatitis to healthy controls. **Methods** Urine was obtained from healthy controls (HC, n=5), inpatients with mild acute pancreatitis (AP, n=5), and outpatients with chronic pancreatitis (CP, n=5). Proton nuclear magnetic resonance spectra were obtained for each sample. Metabolites were identified and quantified in each spectrum; resulting concentrations were normalized to account for differences in dilution among samples. Kruskal-Wallis test, post-hoc Mann-Whitney U tests, and principal component analysis were performed to identify metabolites that discriminate healthy controls, acute pancreatitis, and chronic pancreatitis. **Results** Sixty metabolites were identified and quantified; five were found to differ significantly (P<0.05) among the three groups. Of these, citrate and adenosine remained significant after validation by random permutation. Principal component analysis demonstrated that healthy control urine samples can be differentiated from patients with chronic pancreatitis or acute pancreatitis; chronic pancreatitis patients could not be distinguished from acute pancreatitis patients. **Conclusions** This metabolomic investigation demonstrates that this non-invasive technique offers insight into the metabolic states of pancreatitis. Although the identified metabolites cannot conclusively be defined as biomarkers of disease, future studies will validate our findings in larger patient cohorts.

INTRODUCTION

Diseases of the pancreas affect greater than 1 million persons in the United States annually, resulting in nearly \$3 billion in direct and indirect medical costs. The most prevalent disorder of the pancreas is pancreatitis. This disease is characterized by inflammation of the exocrine portion of the pancreas that secretes digestive enzymes into the duodenum. Pancreatitis can take two forms, acute and chronic. Acute pancreatitis begins abruptly and either resolves or worsens within several days. In contrast, chronic pancreatitis is a chronic inflammatory disorder with a protracted disease course developing and progressing over several years.

Acute pancreatitis is the most common cause of hospitalization for pancreatic disorders in the United States and is associated with significant resource utilization, morbidity and mortality. Recent national survey data indicate increasing frequency of hospitalizations (4.6 of every 1,000 hospital admissions, greater than 200,000 admissions in the U.S. annually) and costs (\$2 billion in direct annual costs) associated with acute pancreatitis in the United States [1, 2]. Currently, diagnoses of acute pancreatitis are made by standard laboratory testing and radiologic imaging [3]. Severe cases may be associated with complications such as organ failure, necrosis, and death; approximately 2% of total acute pancreatitis attacks are fatal.

Although the exact pathogenesis of chronic pancreatitis remains to be determined, a variety of etiologic risk factors have been outlined, including repeated bouts of acute pancreatitis [4]. Objective features of chronic pancreatitis, detectable by current diagnostic radiologic, endoscopic and biochemical studies, are associated with moderate to advanced stage disease. However, at these stages, pathological structural and functional changes are irreversible and only symptomatic treatment is possible [5, 6]. Furthermore, the advanced stages of chronic pancreatitis are associated with diabetes, severe pain, malabsorption of

Received November 13th, 2012 – Accepted November 27th, 2012

Key words Metabolomics; Nuclear Magnetic Resonance, Biomolecular; Pancreatitis, Acute Necrotizing; Pancreatitis, Chronic; Urine

Abbreviations AP: acute pancreatitis; CP: chronic pancreatitis; HC: healthy controls; HMDB: Human Metabolome Database; NMR: nuclear magnetic resonance; PCA: principal component analysis

Correspondence Elizabeth R Lusczek
Critical Care and Acute Care Surgery; University of Minnesota;
516 Delaware St. SE; Minneapolis, MN 55455; USA
Phone: +1-612.624.8661; Fax: +1:612.625.3675
E-mail: lusc0006@umn.edu

nutrition, and the development of pancreatic cancer. The prevalence of chronic pancreatitis in industrialized countries, although frequently misdiagnosed [7] or neglected [8], is approximately 30 per 100,000 inhabitants [9, 10]. No specific therapy is available to arrest the acute inflammatory response seen in acute pancreatitis or to retard the progression of chronic pancreatitis.

Metabolomics, the systematic, high-throughput profiling of the host of metabolites in an organism, can be used to characterize an organism's metabolic response to a pathophysiologic state [11]. Nuclear magnetic resonance (NMR) spectroscopy-based approaches are well-suited for high-throughput studies, offering the ability to profile a wide range of metabolites in a single spectrum with minimal sample preparation [12, 13]. Such studies are commonly used to identify biomarkers or sets of biomarkers of disease [14, 15, 16].

The characterization of the metabolic response in pancreatitis is a potentially useful strategy to determine the effects of pancreatic injury in both chronic and acute pancreatitis. Improved understanding of the metabolic consequences of these diseases may yield non-invasive diagnostic biomarkers and therapeutic targets. To that end, we performed a pilot study to determine if differences could be detected among NMR-based urinary metabolic profiles of patients with chronic pancreatitis, acute pancreatitis, and healthy controls.

METHODS

Study Population

The study population comprised adult patients referred to the Center for Pancreatic Disease at Brigham and Women's Hospital in Boston, MA, USA for evaluation of abdominal pain, and healthy adult volunteers with no pancreatic disease or risk factors for pancreatic disease. Three cohorts were studied: inpatients with mild acute pancreatitis (AP, n=5), outpatients with chronic pancreatitis (CP, n=5), and healthy controls (HC, n=5).

Urine Collection

Clean-catch urine was collected for all subjects. Second void of the day urine was collected using the clean-catch method for acute and chronic pancreatitis patients. Healthy control and chronic pancreatitis collections were done in the outpatient setting. Acute pancreatitis samples were collected within 48 hours of admission. All urine was promptly aliquoted and stored at -80°C until analysis.

Urine Sample Processing

Urine samples were thawed at the time of preparation for NMR analysis. One mL of thawed urine was mixed with 0.5 mL of 0.2 M sodium phosphate buffer prepared with D₂O to control pH. The mixture was

placed on ice for 10 minutes and then centrifuged at 7,000 g for 10 minutes. A total of 500 µL of the supernatant was withdrawn and combined with 50 µL of the internal standard 3-(trimethylsilyl)propionic acid (TSP, Sigma-Aldrich, St. Louis, MO, USA) to a concentration of 1 mM [17]. The internal standard and buffer were prepared with D₂O to provide a lock for the NMR signal. The pH of the final solution was recorded and the mixture was transferred to separate 5 mm NMR tubes (Wilmad, LabGlass, Vineland, NJ, USA).

Nuclear Magnetic Resonance (NMR) Spectroscopy

Proton NMR spectra were collected with a Bruker Avance spectrometer with autosampler and 5 mm triple resonance ¹H/¹³C/¹⁵N TXI CryoProbe with Z-gradient, running TopSpin v. 2.16 (Bruker BioSpin, Fremont, CA, USA) at 700.13 MHz. A 1D NOESY (nuclear Overhauser effect spectroscopy) pulse sequence was used. The 90° pulse width was calibrated for each sample, and was generally 12-13 µs. The relaxation time was defined by each sample's 90° pulse width. The relaxation delay was 2 s, the acquisition time was 3 s, the spectral width was 10 kHz, the total number of data points collected was 63,000, and the number of transients collected was 128, for a total experiment time of 11 minutes and 17 seconds. During the relaxation period, the water resonance was presaturated. All spectra were collected at a temperature of 298 °K. Line broadening at 0.5 Hz was applied before fast Fourier transform (FFT); autophasing and auto-baseline correction were applied by TopSpin.

Chemomx software (Edmonton, AB, Canada) [18] was used to identify and quantify a portion of the metabolites present in each urine sample. Fine manual phasing and baseline corrections and the software's Reference Deconvolution algorithm was applied to each spectrum before targeted profiling of the metabolites was performed. Sixty metabolites were fit in each urine sample in this study, resulting in a profile containing the concentration of each identified metabolite in millimoles per liter (mM). The metabolomic profiles containing the urine concentrations were then normalized using the probabilistic quotient method [19, 20] to correct for differences in dilution among samples.

ETHICS

Informed consent was given by all patients and healthy controls enrolled in this study. This protocol was approved by the Institutional Review Board at Brigham and Women's Hospital (BWH; IRB #2007-P-002480/1). The study protocol conforms to the ethical guidelines of the "World Medical Association (WMA) Declaration of Helsinki - Ethical Principles for Medical Research Involving Human Subjects" adopted by the 18th WMA General Assembly, Helsinki, Finland, June 1964 and amended by the 59th WMA General Assembly, Seoul, South Korea, October 2008.

STATISTICS

Data are reported as mean and range or mean ± standard deviation (SD). Significant urinary metabolites according to the three groups (acute pancreatitis, chronic pancreatitis, healthy control) were identified by Kruskal-Wallis test. Differences between metabolite concentrations were assessed with post-hoc Wilcoxon rank-sum tests with Bonferroni correction to accommodate the small sample size. These results were

then validated with random permutations. All statistical tests were done with R software (R Foundation for Statistical Computing, <http://cran.r-project.org/>) [21]. R software was also used to generate a principal component analysis model with respect to patient group, and to create boxplots of profiled metabolites (Supplementary Figure).

To check for consistency, absolute metabolite concentrations were normalized to creatinine and compared to values reported in the Human

Table 1. Demographics for acute pancreatitis (AP) patients, chronic pancreatitis (CP) patients, and healthy controls (HC). Observed low peak HCO₃⁻ concentration values and Cambridge III imaging scores for chronic pancreatitis patients were consistent with advanced disease.

ID	Gender	Race	Age (years)	MANNHEIM CP diagnosis	Atlanta AP severity	Ranson criteria	Imaging	Etiology	TIGAR-O CP classification
Acute pancreatitis									
AP1	Female	White	58	N/A	Mild	2	Normal MRI	Gallstone	N/A
AP2	Male	Black	42	N/A	Mild	2	Balthazar C	Alcohol	N/A
AP3	Male	White	44	N/A	Mild	1	Balthazar C	Obstructive	N/A
AP4	Male	Not known	59	N/A	Mild	2	Diffuse enlargement on abdominal US	Alcohol	N/A
AP5	Female	Black	63	N/A	Mild	2	Balthazar A	Gallstone	N/A
Chronic pancreatitis									
CP1	Male	White	66	Definite	N/A	N/A	Cambridge III	Alcohol, smoking	Toxic-metabolic
CP2	Female	White	52	Definite	N/A	N/A	Cambridge III	Alcohol, smoking, RAP	Toxic-metabolic
CP3	Male	White	77	Definite	N/A	N/A	Cambridge III	Alcohol, smoking	Toxic-metabolic
CP4	Male	White	42	Definite	N/A	N/A	Cambridge III	Alcohol, hyperlipidemia	Toxic-metabolic
CP5	Female	White	62	Definite	N/A	N/A	Cambridge III	Alcohol, smoking	Toxic-metabolic
Healthy controls									
HC1	Male	White	24	N/A	N/A	N/A	N/A	N/A	N/A
HC2	Male	Asian	32	N/A	N/A	N/A	N/A	N/A	N/A
HC3	Female	White	34	N/A	N/A	N/A	N/A	N/A	N/A
HC4	Female	White	62	N/A	N/A	N/A	N/A	N/A	N/A
HC5	Female	White	33	N/A	N/A	N/A	N/A	N/A	N/A

RAP: recurrent acute pancreatitis

Table 1. Continued.

ID	Peak HCO ₃ ⁻ concentration (mEq/L)	Smoking status	Smoking amount (packs/year)	Alcohol consumption	Alcohol amount
Acute pancreatitis					
AP1	N/A	Current	20	Not known	Not known
AP2	N/A	Never	0	Current	Moderate
AP3	N/A	Past	17	Past	Excessive
AP4	N/A	Current	>30	Current	Excessive
AP5	N/A	Current	67.5	Not known	Not known
Chronic pancreatitis					
CP1	39	Current	96	Current	Increased
CP2	38	Current	10	Past	Excessive
CP3	36	Past	25	Past	Excessive
CP4	40	Never	0	Current	Increased
CP5	37	Past	>40	Past	Excessive
Healthy controls					
HC1	N/A	N/A	N/A	N/A	N/A
HC2	N/A	N/A	N/A	N/A	N/A
HC3	N/A	N/A	N/A	N/A	N/A
HC4	N/A	N/A	N/A	N/A	N/A
HC5	N/A	N/A	N/A	N/A	N/A

Table 2. Comparison of select metabolite concentrations obtained from the HMDB (Human Metabolome Database) and from these experiments. For the sake of comparison, absolute metabolite concentrations were normalized to creatinine concentrations. Values are reported as mean (range) with units of $\mu\text{mol}/\text{mmol}$ creatinine.

Common urinary metabolites	HMDB healthy reference ^a	Acute pancreatitis	Chronic pancreatitis	Healthy controls
Adenosine	1.4 (0.9-2.3)	3.6 (1.3-5.3)	4.4 (3.1-6.4)	1.6 (0.31-3.9)
Alanine	21.8 (7.1-43.1)	20 (10-41)	15 (10-29)	19 (12-33)
Choline	3.5 (1.4-6.1)	11 (3.6-19)	7.9 (2.5-18)	3.1 (1.3-4.9)
Citrate	203 (49-600)	120 (6.6-260)	82 (10-250)	240 (130-340)
Formate	26.8 (6.0-120.9)	33 (8.1-63)	12 (2.2-24)	27 (15-49)
Hippurate	198 (26-718)	170 (39-320)	120 (14-250)	120 (59-205)
Lactate	11.6 (3.5-29.3)	15 (13-22)	12 (7.7-16)	17 (11-28)
Quinolate	5.2 (1.0-17.5)	6.4 (4.5-9.1)	5.2 (2.0-12)	7.7 (2.1-17)
Trigonelline	31.1 (5.5-109.3)	16 (1.8-40)	27 (0.60-81)	29 (3.2-73)
Valine	5.5 (2.7-9.8)	4.1 (2.1-6.1)	3.4 (1.7-6.3)	3.8 (2.9-5.0)

^a HMDB healthy reference values were obtained via NMR as reported by S. Bouatra *et al.* 2012 (The Human Urine Metabolome; manuscript in preparation: <http://www.hmdb.ca>)

Metabolome Database (HMDB; <http://www.hmdb.ca>) [22]. Values obtained herein were compared to those also obtained by NMR as reported in HMDB entries (S. Bouatra *et al.*, 2012, The Human Urine Metabolome; manuscript in preparation: <http://www.hmdb.ca>).

RESULTS

Patient characteristics are listed in Table 1. All of the patients with chronic pancreatitis had alcohol consumption as a related factor and half of the patients with acute pancreatitis had alcohol consumption as a related factor. Peak HCO_3^- concentrations were obtained from i.v. secretin-stimulated endoscopic pancreatic function tests (ePFT) [23]. Accepted normal values are greater than 75 mEq/L [24]. The observed low peak HCO_3^- concentration values and Cambridge III imaging scores for chronic pancreatitis patients were consistent with advanced disease.

Nuclear Magnetic Resonance (NMR) Spectroscopy Identified Metabolites in the Urine of all Three Cohorts

Sixty metabolites were identified and quantified in the urine. Urea was subsequently omitted from the data set because its protons are in exchange with water and the signal is compromised by water suppression. Boxplots of the remaining 59 metabolites and their averages and standard deviations are reported in the Supplementary Table and Figure. Concentrations of a subset of metabolites identified here are compared with those reported in the Human Metabolome Database in Table 2 and demonstrate that the results of this study are comparable to known concentration ranges.

Of the 59 metabolites analyzed in this study, five were found to differ significantly among the three groups by Kruskal-Wallis test: acetone, adenosine, citrate, ribose, and 3-indoxylsulfate. These five metabolites were then subjected to pairwise Wilcoxon rank-sum tests with Bonferroni correction in a post-hoc evaluation (Table 3). Validation by random permutation confirmed that of these, only citrate and adenosine were significantly

different among the groups. Citrate and adenosine concentrations in each of the groups are shown in Figure 1. Urinary citrate concentrations in healthy controls were greater than those in chronic pancreatitis patients ($P=0.019$, validation by random permutation). Conversely, urinary adenosine concentrations were lower in healthy controls than in chronic pancreatitis patients ($P=0.026$, validation by random permutation).

Principal Component Analysis Revealed Separation Between Healthy Controls and Pancreatitis Patients

Principal component analysis (PCA) was used to model the metabolic profiles according to group (chronic pancreatitis, acute pancreatitis, or healthy control). The first three components collectively explain 41% of the variance in the data (PC1=16%, PC2=13%, PC3=12%). Examination of the three-dimensional scores plot (Figure 2) showed that the control urine samples in green were generally separated from the urine samples obtained from pancreatitis patients. PCA demonstrated no separation between chronic

Table 3. Significant urinary metabolites identified by Kruskal-Wallis test according to group (AP: acute pancreatitis, CP: chronic pancreatitis, HC: healthy control). Wilcoxon rank-sum tests with Bonferroni correction are then applied to test these metabolites; those with significant P values ($P<0.05$) are bolded. After the significant Wilcoxon rank-sum test results are validated by random permutation, only citrate and adenosine results are significant ($P=0.019$ and $P=0.026$, respectively).

Significant metabolites ^a	Exact P value ^b	Comparison by group
Acetone	0.095	AP > CP
Acetone	0.024	AP > HC
Adenosine	0.048	CP > HC
Citrate	0.095	AP < HC
Citrate	0.048	CP < HC
Ribose	0.048	AP > HC
Ribose	0.095	CP > HC
3-Indoxylsulfate	0.095	AP < HC

^a Identified by Kruskal-Wallis test

^b Wilcoxon rank-sum test with Bonferroni correction




Figure 1. Boxplots depict concentrations of two significant metabolites identified after Wilcoxon rank-sum tests and validation by random permutation. Data are reported in $\mu\text{mol/L}$ (normalized). Citrate (left) concentration was significantly lower in chronic pancreatitis (CP) patients compared to healthy controls (* $P=0.019$). Adenosine (right) concentration was significantly higher in chronic pancreatitis patients compared to healthy controls (** $P=0.026$). The box covers the first (Q1) and third (Q3) quartile of the data. The line in the box represents the median value. The whiskers extend to ± 1.5 interquartile range (IQR), where $\text{IQR}=\text{Q3}-\text{Q1}$. Outliers are shown as points.

pancreatitis and acute pancreatitis urine samples. The loadings plot (Figure 3) demonstrated the contribution of each of the metabolites to the model; citrate and adenosine are highlighted in blue. Though there is little pattern in the distribution of the metabolites, it is clear




Figure 2. Principal component (PC) scores plot of urinary metabolic profiles shows acute pancreatitis (AP: black), chronic pancreatitis (CP: red), and healthy control (HC: green) urine samples in three dimensions. Separation of the healthy control urine samples from acute pancreatitis and chronic pancreatitis samples is observed. No separation in acute pancreatitis and chronic pancreatitis urine samples is observed.

that citrate and adenosine lie far from each other in the loadings.

DISCUSSION

This pilot study demonstrates that the urinary metabolome can be used to differentiate patients with acute or chronic pancreatitis from healthy controls. Although only two potential biomarkers were identified, establishing the utility of this methodology is important for future metabolomic studies of pancreatic disease.

Of five identified metabolites that differed significantly among groups by Kruskal-Wallis test, four remained significant after evaluation by Wilcoxon rank-sum tests with Bonferroni correction (acetone, adenosine, citrate, ribose). Of these metabolites, only adenosine and citrate survived validation by random permutation. These two metabolites could reasonably be expected to differ among pancreatitis patients and healthy controls. Citrate is known to be a significant component of pancreatic secretions [25]. Given that alcohol use or abuse has been associated with a decrease in citrate in pancreatic secretions [26], it is notable that most patients in the chronic pancreatitis and acute pancreatitis groups reported some form of regular alcohol consumption (Table 1). The decreased urinary citrate in the pancreatitis cohorts compared to healthy subjects may be associated with decreased citrate in pancreatic secretions due to alcohol metabolism. This weakens citrate’s candidacy as a biomarker of pancreatitis considerably, as the observed differences are potentially confounded by alcohol consumption. Still, future studies controlling for alcohol consumption may support citrate as a urinary biomarker of pancreatitis. Other etiologies of low urinary citrate




Figure 3. Principal component (PC) loadings plot of urinary metabolic profiles shows the influence of the metabolites on the scores. It can be seen that citrate and adenosine (boxed) are well-separated in the model.

include metabolic acidosis, potassium depletion, and acetazolamide [27, 28]. These conditions do not appear to be relevant in the population studied here.

Adenosine may prove to be a better marker of pancreas disease since it possesses known anti-inflammatory properties [29]. However, adenosine as a marker of inflammation is unlikely to be specific to pancreatitis [30], thus limiting its utility as a biomarker if not validated in further studies. While urinary adenosine concentrations may vary with changes in extracellular adenosine [31], variation in urinary adenosine concentration is known to be low in humans, and is unaffected by sodium content or changes in fluid homeostasis [32]. This implies that renal handling of adenosine does not further confound our results.

As suggested above, this study has several limitations, which may be attributed to small sample size. We attempted to minimize the effects of sample size on the statistical analysis of the data by using Kruskal-Wallis global test, followed by Wilcoxon rank-sum tests with Bonferroni correction and validation by random permutation. We also chose to use PCA, an unsupervised multivariate analysis technique, instead of a supervised technique. Supervised analyses such as partial least squares discriminant analysis (PLS-DA) are common in large metabolomics studies, but are inappropriate for small studies such as this. Additionally, we have not accounted for all confounding variables. As discussed, citrate's relationship to alcohol consumption compromises its usefulness as a biomarker of pancreatitis. Other factors such as age, differences in liver function, and dietary status likely exist among the patients providing samples. Furthermore, the possibility of differences among patient populations, etiology, and disease severity inducing variations in metabolomes presents an additional challenge. A larger study of the urine, serum, and pancreas fluid metabolomes with well-defined patient enrollment criteria, controlling for age, disease severity, and etiology of pancreatitis will account for confounding factors. This would result in a higher proportion of variance explained in the PCA model. A larger study with better-defined cohorts would also allow for partial least squares discriminant analysis modeling. We are currently performing a larger metabolomics study of ERCP-induced pancreatitis that will address many of these issues.

Our study demonstrates that metabolomics techniques are capable of distinguishing urine samples obtained from pancreatitis patients and from those obtained from healthy controls. This is a novel, non-invasive technique that can provide insight into the metabolic states of patients with chronic and acute pancreatitis. The significant increase in urinary adenosine and decrease in urinary citrate among pancreatitis patients compared to healthy controls may have reflected the patients' inflammatory state and alcohol consumption, and as such, validation studies will have to be performed. Although we cannot definitively conclude that these metabolites are biomarkers of pancreatitis,

further analyses with larger cohorts will refine our results and define their diagnostic utility. The methodology described here presents a definitive strategy upon which further analyses with larger cohorts can be tested.

Financial support None

Conflicts of interest None

References

1. Fagenholz PJ, Castillo CFD, Harris NS, Pelletier AJ, Camargo CA. Increasing United States hospital admissions for acute pancreatitis, 1988–2003. *Ann Epidemiol.* 2007; 17(7):491-497. [PMID: 1744682]
2. Fagenholz PJ, Fernández-del Castillo C, Harris NS, Pelletier AJ, Camargo CA. National study of united states emergency department visits for acute pancreatitis, 1993–2003. *BMC emergency medicine.* 2007; 7(1):1. [PMID: 17241461]
3. Cappell MS. Acute pancreatitis: Etiology, clinical presentation, diagnosis, and therapy. *Med Clin North Am.* 2008; 92(4):889-923. [PMID: 18570947]
4. Etemad B, Whitcomb DC. Chronic pancreatitis: Diagnosis, classification, and new genetic developments. *Gastroenterology* 2001; 120(3):682-707. [PMID: 1179244]
5. Warshaw AL, Banks PA, Fernández-del Castillo C. AGA technical review: Treatment of pain in chronic pancreatitis. *Gastroenterology* 1998; 115(3):765-776. [PMID: 9721175]
6. Witt H, Apte MV, Keim V, Wilson JS. Chronic pancreatitis: Challenges and advances in pathogenesis, genetics, diagnosis, and therapy. *Gastroenterology* 2007; 132(4):1557-1573. [PMID: 17466744]
7. Flasar MH, Goldberg E. Acute abdominal pain. *Med Clin North Am* 2006; 90(3):481-504. [PMID: 16473101]
8. Lankisch P. The problem of diagnosing chronic pancreatitis. *Dig Liver Dis* 2003; 35(3):131-134. [PMID: 12779064]
9. Otsuki M. Chronic pancreatitis in japan: Epidemiology, prognosis, diagnostic criteria, and future problems. *J Gastroenterol* 2003; 38(4):315-326. [PMID: 127437700]
10. Whitcomb DC, Yadav D, Adam S, et al. Multicenter approach to recurrent acute and chronic pancreatitis in the United States: The North American pancreatitis study 2 (NAPS2). *Pancreatol* 2008; 8(4-5):520-531. [PMID: 18765957]
11. Fiehn O. Metabolomics—the link between genotypes and phenotypes. *Plant Mol Biol* 2002; 48(1):155-171. [PMID: 11860207]
12. Nicholson JK, Lindon JC, Holmes E. 'Metabonomics': Understanding the metabolic responses of living systems to pathophysiological stimuli via multivariate statistical analysis of biological NMR spectroscopic data. *Xenobiotica* 1999; 29(11):1181-1189. [PMID: 10598751]
13. Madsen R, Lundstedt T, Trygg J. Chemometrics in metabolomics—A review in human disease diagnosis. *Anal Chim Acta.* 2010; 659(1-2):23-33. [PMID: 20103103]
14. Sreekumar A, Poisson LM, Rajendiran TM, et al. Metabolomic profiles delineate potential role for sarcosine in prostate cancer progression. *Nature* 2009; 457(7231):910-914. [PMID: 19212411]
15. Spratlin JL, Serkova NJ, Eckhardt SG. Clinical applications of metabolomics in oncology: A review. *Clin Cancer Res* 2009; 15(2):431-40. [PMID: 19147747]
16. Carraro S, Rezzi S, Reniero F, et al. Metabolomics applied to exhaled breath condensate in childhood asthma. *Am J Respir Crit Care Med* 2007; 175(10):986-90. [PMID: 17303796]
17. Mortishire-Smith RJ, Skiles GL, Lawrence JW, et al. Use of metabolomics to identify impaired fatty acid metabolism as the mechanism of a drug-induced toxicity. *Chem Res Toxicol* 2004; 17(2):165-173. [PMID: 14967004]

18. Weljie AM, Newton J, Mercier P, Carlson E, Slupsky CM. Targeted profiling: Quantitative analysis of 1H NMR metabolomics data. *Anal Chem* 2006; 78(13):4430-4442. [PMID: 16808451]
19. Dieterle F, Ross A, Schlotterbeck G, Senn H. Probabilistic quotient normalization as robust method to account for dilution of complex biological mixtures. application in 1H NMR metabonomics. *Anal Chem* 2006; 78(13):4281-4290. [PMID: 16808434]
20. Luszczek ER, Nelson T, Lexcen D, Witowski NE, Mulier KE, Beilman G. Urine metabolomics in hemorrhagic shock: Normalization of urine in the face of changing intravascular fluid volume and perturbations in metabolism. *J Bioanal Biomed*. 2011; 3:038-048.
21. R Development Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing Vienna Austria. 2010; <http://cran.r-project.org/>.
22. Wishart DS, Knox C, Guo AC, et al. HMDB: A knowledgebase for the human metabolome. *Nucleic Acids Res*. 2009; 37(suppl 1):D603-D610. [PMID: 18953024]
23. Wu B, Conwell DL. The endoscopic pancreatic function test. *Am J Gastroenterol*. 2009; 104(10):2381-2383. [PMID: 19806083]
24. Stevens T, Conwell DL, Zuccaro G, Lewis SA, Love TE. The efficiency of endoscopic pancreatic function testing is optimized using duodenal aspirates at 30 and 45 minutes after intravenous secretin. *Am J Gastroenterol*. 2007; 102(2):297-301. [PMID: 17100964]
25. Boustière C, Sarles H, Lohse J, Durbec J, Sahel J. Citrate and calcium secretion in the pure human pancreatic juice of alcoholic and nonalcoholic men and of chronic pancreatitis patients. *Digestion* 1985; 32(1):1-9. [PMID: 4018438]
26. Sarles H, Bernard J, Johnson C. Pathogenesis and epidemiology of chronic pancreatitis. *Annu Rev Med* 1989; 40(1):453-468. [PMID: 2658760]
27. Simpson D. Citrate excretion: A window on renal metabolism. *American Journal of Physiology-Renal Physiology*. 1983; 244(3):F223-F234. [PMID: 6338740]
28. Hamm LL. Renal handling of citrate. *Kidney Int*. 1990; 38(4):728-735. [PMID: 2332510]
29. Noji T, Nan-ya K, Mizutani M, et al. KF24345, an adenosine uptake inhibitor, ameliorates the severity and mortality of lethal acute pancreatitis via endogenous adenosine in mice. *Eur J Pharmacol* 2002; 454(1):85-93. [PMID: 12409009]
30. Bours M, Swennen E, Di Virgilio F, Cronstein B, Dagnelie P. Adenosine 5'-triphosphate and adenosine as endogenous signaling molecules in immunity and inflammation. *Pharmacol Ther* 2006; 112(2):358-404. [PMID: 1678477]
31. Thompson CI, Sparks HV, Spielman WS. Renal handling and production of plasma and urinary adenosine. *American Journal of Physiology-Renal Physiology*. 1985; 248(4):F545-F551. [PMID: 3985162]
32. Heyne N, Benöhr P, Mühlbauer B, Delabar U, Risler T, Osswald H. Regulation of renal adenosine excretion in humans—role of sodium and fluid homeostasis. *Nephrology Dialysis Transplantation*. 2004; 19(11):2737-2741. [PMID: 15385638]

Supplementary Table. Absolute (non-normalized) concentrations of 59 metabolites profiled by NMR in each urine sample for acute pancreatitis patients, chronic pancreatitis patients, and healthy controls. Values are reported as mean±SD with units of µmol/L. Boxplots of each metabolite (non-normalized concentrations) are shown the Supplementary Figure.

Metabolite	Acute pancreatitis	Chronic pancreatitis	Healthy controls	Metabolite	Acute pancreatitis	Chronic pancreatitis	Healthy controls
1,6-Anhydro β D glucose	70±86	65±37	170±220	Kynurenine	84±141	89±37	24±4.7
1-Methylnicotinamide	7.1±2.9	230±230	20±21	Lactate	32±15	69±39	66±54
2-Aminoadipate	150±120	450±340	140±47	Lactose	66±59	130±72	70±46
2-Methylglutarate	13±9.6	18±7.4	24±13	Malonate	68±38	290±360	89±41
2-Oxoglutarate	34±24	95±70	58±49	Mannose	31±32	23±17	8.9±7.9
3-Hydroxyisovalerate	8.1±4.1	21±10	17±9.0	Methylguanidine	11±12	47±78	18±16
3-Indoxylsulfate	32±30	162±110	100±59	N,N Dimethylglycine	8.7±3.3	42±45	11±5.0
4-Aminohippurate	16±5.3	110±140	52±55	N-Methylhydantoin	11±2.8	40±27	20±8.8
ADP	3.1±1.9	10±8.3	5.4±5.7	N-Phenylacetyl glycine	100±49	330±240	140±69
Acetate	16±7.9	45±24	29±15	O-Acetylcholine	23±22	42±50	24±16
Acetoacetate	24±24	130±109	35±12	O-Phosphocholine	32±24	43±37	15±6.9
Acetone	84±100	32±22	9.5±4.1	Phenylacetyl glycine	83±80	400±330	98±82
Adenosine	7.3±4.5	28±19	4.8±3.6	Pyruvate	11±6.4	34±25	14±3.4
Alanine	35±7.7	86±68	68±46	Quinolate	15±11	28±24	27±25
Allantoin	39±26	67±31	30±11	Ribose	230±220	440±290	82±80
Betaine	30±30	420±650	26±18	Succinate	10±11	41±21	35±26
Caffeine	31±18	67±25	43±25	Taurine	230±200	380±260	120±57
Choline	24±22	45±33	11±5.9	Threonine	33±14	98±100	63±30
Citrate	180±110	430±630	910±620	Trigonelline	28±26	83±84	89±60
Creatine	51±39	170±88	89±69	Trimethylamine	2.5±1.7	12±8.7	4.6±2.4
Creatinine	2,200±1,260	6,100±3,800	3,500±1,600	Trimethylamine N-oxide	110±100	520±690	140±110
Dimethylamine	100±75	220±74	120±48	Tryptophan	15±7.1	41±34	22±19
Ethanolamine	76±40	200±140	180±130	Tyramine	21±15	72±33	36±20
Formate	52±31	57±39	96±67	Tyrosine	14±6.9	40±21	27±18
Fumarate	2.9±1.7	3.9±2.2	2.7±1.3	Valine	7.3±1.8	19±12	13±6.8
Glucose	8,200±16,820 ^a	420±430	110±190	Xanthine	49±14	97±88	38±22
Glycine	130±63	370±500	300±191	cis-Aconitate	80±58	200±120	130±59
Hippurate	340±290	530±480	410±270	pi.Methylhistidine	84±36	440±590	100±60
Hypoxanthine	9.2±6.8	97±181	14±5.2	tau.Methylhistidine	56±43	120±91	48±18
Inosine	10±7.5	25±33	5.2±4.9				

^a One acute pancreatitis patient had a raw glucose of 38 mM. Without this value, mean glucose values for acute pancreatitis patients are 650±110 µmol/L.

Supplementary Figure. Boxplots of absolute (non-normalized) concentrations of 59 profiled urinary metabolites. The box covers the first (Q1) and third (Q3) quartile of the data. The line in the box represents the median value. The whiskers extend to ± 1.5 interquartile range (IQR), where $IQR=Q3-Q1$. Outliers are shown as points. Concentration units are expressed in $\mu\text{mol/L}$.

